

Polymicrobial Nature of Tick-Borne Diseases

Santiago Sanchez-Vicente,^a Teresa Tagliaferro,^b James L. Coleman,^a Jorge L. Benach,^a Rafal Tokarz^b

^aDepartment of Molecular Genetics and Microbiology and Center for Infectious Diseases, Stony Brook University, Stony Brook, New York, USA

^bCenter for Infection and Immunity, Mailman School of Public Health, Columbia University, New York, New York, USA

ABSTRACT Tick-borne diseases have doubled in the last 12 years, and their geographic distribution has spread as well. The clinical spectrum of tick-borne diseases can range from asymptomatic to fatal infections, with a disproportionate incidence in children and the elderly. In the last few years, new agents have been discovered, and genetic changes have helped in the spread of pathogens and ticks. Polymicrobial infections, mostly in *Ixodes scapularis*, can complicate diagnostics and augment disease severity. *Amblyomma americanum* ticks have expanded their range, resulting in a dynamic and complex situation, possibly fueled by climate change. To document these changes, using molecular biology strategies for pathogen detection, an assessment of 12 microbes (9 pathogens and 3 symbionts) in three species of ticks was done in Suffolk County, New York. At least one agent was detected in 63% of *I. scapularis* ticks. *Borrelia burgdorferi* was the most prevalent pathogen (57% in adults; 27% in nymphs), followed by *Babesia microti* (14% in adults; 15% in nymphs), *Anaplasma phagocytophilum* (14% in adults; 2% in nymphs), *Borrelia miyamotoi* (3% in adults), and Powassan virus (2% in adults). Polymicrobial infections were detected in 22% of *I. scapularis* ticks, with coinfections of *B. burgdorferi* and *B. microti* (9%) and of *B. burgdorferi* and *A. phagocytophilum* (7%). Three *Ehrlichia* species were detected in 4% of *A. americanum* ticks. The rickettsiae constituted the largest prokaryotic biomass of all the ticks tested and included *Rickettsia amblyommatis*, *Rickettsia buchneri*, and *Rickettsia montanensis*. The high rates of polymicrobial infection in ticks present an opportunity to study the biological interrelationships of pathogens and their vectors.

IMPORTANCE Tick-borne diseases have increased in prevalence in the United States and abroad. The reasons for these increases are multifactorial, but climate change is likely to be a major factor. One of the main features of the increase is the geographic expansion of tick vectors, notably *Amblyomma americanum*, which has brought new pathogens to new areas. The clinical spectrum of tick-borne diseases can range from asymptomatic to fatal infections, with a disproportionate incidence in children and the elderly. In addition, new pathogens that are cotransmitted by *Ixodes scapularis* have been discovered and have led to difficult diagnoses and to disease severity. Of these, *Borrelia burgdorferi*, the agent of Lyme disease, continues to be the most frequently transmitted pathogen. However, *Babesia microti*, *Borrelia miyamotoi* (another spirochete), *Anaplasma phagocytophilum*, and Powassan virus are frequent cotransmitted agents. Polymicrobial infection has important consequences for the diagnosis and management of tick-borne diseases.

KEYWORDS *Amblyomma*, *Anaplasma*, *Babesia*, *Borrelia burgdorferi*, *Ehrlichia*, *Ixodes*, Lyme disease, Powassan, *Rickettsia*

Polymicrobial infections are known for a variety of disorders, such as urinary tract infections, sexually transmitted diseases, periodontal disease, otitis media, and opportunistic infections associated with hospital procedures, to name just a few. In

Citation Sanchez-Vicente S, Tagliaferro T, Coleman JL, Benach JL, Tokarz R. 2019. Polymicrobial nature of tick-borne diseases. mBio 10:e02055-19. <https://doi.org/10.1128/mBio.02055-19>.

Editor Liise-anne Pirofski, Albert Einstein College of Medicine

Copyright © 2019 Sanchez-Vicente et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Jorge L. Benach, Jorge.benach@stonybrook.edu, or Rafal Tokarz, Rt2249@cumc.columbia.edu.

This article is a direct contribution from a Fellow of the American Academy of Microbiology. Solicited external reviewers: Abdu Azad, University of Maryland School of Medicine; Erol Fikrig, Yale School of Medicine; Ulrike Munderloh, University of Minnesota; Sam R. Telford III, Tufts University School of Veterinary Medicine.

Received 5 August 2019

Accepted 7 August 2019

Published 10 September 2019

these settings, coexisting microbes can generate synergy or interference of the infectious process. However, regardless of the outcome of a specific polymicrobial infection, the interrelationships among the microbes are likely to have an impact on the course of the infection. Unlike some of the examples above where polymicrobial infections affect only the patient, the various pathogens and symbionts of ticks interact in both the tick itself and in the patient with consequences that have not been totally foreseen (1).

Tick-borne diseases have become a worldwide threat to public health. In the United States, cases increased from 22,000 in 2004 to >48,000 in 2016 (2). Tick-borne diseases range from subclinical to fatal infections, with a disproportionate incidence in children and the elderly. Moreover, some can also be transmitted by blood transfusions and cause severe disease in patients with underlying disorders. New agents have also been discovered (3, 4). Notably, formerly geographically confined tick species have expanded their range, resulting in a dynamic and complex situation, possibly fueled by climate change. Polymicrobial infections represent another aspect of tick-borne diseases that can complicate diagnosis and augment disease severity. Since the discovery of *Borrelia burgdorferi* in 1982 (5) and human babesiosis caused by the hemoprotozoan *Babesia microti* (6, 7) in Long Island and Nantucket in the late 1970s, more cotransmitted infections have been recognized. Two early serosurveys disclosed a link between babesiosis and erythema migrans and double infections (8, 9). Since then, *Ixodes scapularis* has been implicated as the vector of five human pathogens in the northeast United States. Polymicrobial infections occur in both North America and Europe (10).

Over the past 3 decades, there has been a steady increase in the number of newly discovered tick-borne agents. In addition to *B. burgdorferi* and *B. microti*, *Anaplasma phagocytophilum* (11–15), which was originally classified as a granulocytic *Ehrlichia* species (16), a relapsing fever-like *Borrelia* species, *Borrelia miyamotoi* (3), and the deer tick virus, a variant of Powassan virus (POWV) (4), are also transmitted by *I. scapularis*. POWV was first isolated from a patient with encephalitis (17), and over 100 cases have been reported in the United States (18). The capacity of *I. scapularis* to harbor such a diverse pathogen microbiome increases the risk of polymicrobial infections from a single tick bite. Almost 25 years ago, Telford et al. warned that cotransmission of pathogens will have a unique impact on public health in sites of endemicity (19).

Pathogens transmitted by *Amblyomma americanum* and *Dermacentor variabilis* contribute to the broad spectrum of tick-borne diseases. *A. americanum*, the lone star tick, has historically been found in the southern United States, but it has substantially expanded its range. Anecdotally, it appears that this aggressive species has become the most abundant tick in Long Island, New York, and in other regions previously outside its range (20, 21), and it has done so in a very short time (22, 23). A collateral effect of the expansion of *A. americanum* could be the displacement of *I. scapularis* and *D. variabilis* through competitive interactions that are not understood. Significant shifts in disease prevalence in the future could be due to shifts in the vector populations, and systematic tick-pathogen surveys may answer this question. *A. americanum* is a vector of *Ehrlichia chaffeensis* and *Ehrlichia ewingii* (24–27), both capable of causing severe disease in patients who are elderly or immunodeficient (28). *A. americanum* has been linked to a Lyme disease-like syndrome called southern tick-associated rash illness (STARI). *Borrelia lonestari*, a relapsing fever-like species found in *A. americanum* (29), has been associated with STARI, but its role has not been corroborated. Most recently, *A. americanum* has been implicated in meat allergy syndrome, an intriguing condition that may be due to this tick directly without the intervention of a microbe (30). *Haemaphysalis longicornis*, a newly discovered exotic tick species introduced in the United States (31), is an important vector of human and animal disease agents in its original geographic range (32, 33).

Of note is the increasing awareness of the rickettsial biome in the three vector ticks. The American dog tick, *D. variabilis*, is a vector of *Rickettsia rickettsii*, the agent of Rocky Mountain spotted fever (RMSF). The number of reported cases of spotted fever rickettsioses (SFR) in the United States has increased from 1,713 in 2004 to




FIG 1 Map of Suffolk County, New York, showing locations of tick collection in the northern and southern regions.

4,269 in 2016 (2). However, tick surveillance studies have rarely reported *R. rickettsii* in *D. variabilis* (34, 35). This suggests that other species of *Rickettsia* contribute to the rise in incidence of SFR.

Rickettsia buchneri, an ovarian symbiont, (36), is the most abundant prokaryote in *I. scapularis* (37–39). *Rickettsia amblyommatis* (40) is pervasively associated with *A. americanum* (41). *Rickettsia montanensis* in *D. variabilis* has been implicated circumstantially in an RMSF-like infection (42). These three rickettsial species phylogenetically belong to the spotted fever group but are not agents of disease. However, these rickettsiae, by virtue of their abundance, may have a critical role in pathogen-vector interactions.

The presence of known pathogens as well as probable pathogens and symbionts in ticks requires a polymicrobial approach to the clinical aspects of tick-borne infections. Concurrent polymicrobial infections in humans can have a synergistic effect and result in a more severe course of illness. In addition, the clinical course of one tick-borne disease could be influenced simply by exposure to another microbe. For example, *A. phagocytophilum* and *Ehrlichia* spp. target cells of the innate immune system, and it is possible that even a self-limited transient exposure may influence development of diseases caused by the other pathogens. Excellent reviews have considered the pathogenesis of these organisms (43–46). Of equal importance as the role of polymicrobial infections in clinical disease, the combination of pathogens and symbionts no doubt constitutes a species-specific tick microbiome. There is evidence for detrimental, beneficial, or neutral effects among the interactions of the prokaryotes with each other and with the host tick. These effects can influence access to nutrients, which in turn influence overall fitness of the organisms. For these reasons, and to verify the changing conditions of tick-borne diseases, we performed a polymicrobial assessment of the three species of ticks associated with human disease, including one that is a recent invader. Our results reveal a complex pattern of tick infections with new and emergent pathogens on a background of rapidly shifting tick populations that justify a polymicrobial approach to the study of tick-borne diseases.

RESULTS

Ticks were collected in the spring and fall of 2018 from multiple locations throughout Suffolk, a suburban county that occupies the central and eastern part of Long Island, NY (<https://gisportal.suffolkcountyny.gov/gis/home/>). All locations were clustered into northern sites and southern sites (Fig. 1). In the spring, we collected both

TABLE 1 Prevalence of five pathogens and *R. buchneri* in *I. scapularis* ticks by season^a

Agent ^b	No. of positive ticks (% [95% CI])		
	Spring (n = 198 [140 A/58 N])	Fall (n = 480)	Total (n = 678 [620 A/58 N])
<i>Borrelia burgdorferi</i>			
A	85 (61 [52–69])	266 (55 [51–60])	351 (57 [53–60])
N	16 (27 [17–41])		16 (27 [17–41])
<i>Borrelia miyamotoi</i>			
A	4 (3 [1–8])	16 (3 [2–5])	20 (3 [2–5])
N	0 (0 [0–8])		0 (0 [0–8])
<i>Anaplasma phagocytophilum</i>			
A	19 (14 [9–21])	70 (15 [12–18])	89 (14 [12–17])
N	1 (2 [0.1–10])		1 (2 [0.1–10])
<i>Babesia microti</i>			
A	27 (19 [13–27])	61 (13 [10–16])	88 (14 [12–17])
N	9 (15 [8–28])		9 (15 [8–28])
POWV			
A	5 (3 [1–8])	6 (1 [0.5–3])	11 (2 [1–3])
N	0 (0 [0–8])		0 (0 [0–8])
<i>Rickettsia buchneri</i> ^c			
A	26 (81 [63–92])	64 (89 [79–95])	90 (86 [78–92])
N	21 (91 [70–98])		21 (91 [70–98])

^aTotal prevalences for both spring and fall seasons are included (excluding *R. buchneri*). Values are means and 95% confidence intervals (CI).

^bA, adults; N, nymphs.

^cThirty-two adults and 23 nymphs were tested for the spring season; 72 adult ticks were tested for the fall season.

adults and nymphs of *A. americanum* (676) and *I. scapularis* (198) and adults of *D. variabilis* (296). Fall collections were comprised of 480 *I. scapularis* adults. In total, we examined 1,633 individual ticks and 17 pools of 170 *A. americanum* nymphs (10 nymphs per pool). *I. scapularis* ticks were screened for the presence of *B. burgdorferi*, *B. miyamotoi*, *A. phagocytophilum*, *B. microti*, POWV, and *Rickettsia* spp. We did not screen *I. scapularis* ticks for other ehrlichial agents, because they were not detected in a previous study in our area and their geographic range appears to be limited to the Upper Midwest (39). *A. americanum* ticks were screened for *B. lonestari*, *Ehrlichia* spp., and *Rickettsia* spp., as well as for the five agents tested in *I. scapularis*. *D. variabilis* ticks were screened for *Rickettsia* spp. only.

Pathogen burden of *I. scapularis*. In the northeastern United States, *I. scapularis* is implicated in the transmission of five agents, including *B. burgdorferi*, *A. phagocytophilum*, *B. microti*, *B. miyamotoi*, and POWV. All five agents and *R. buchneri* were detected in our study. A total of 430 of 678 (63%) *I. scapularis* ticks were positive for at least one human pathogen. *B. burgdorferi* was the most prevalent pathogen in both adults and nymphs (57% and 27%, respectively), followed by *B. microti* (14% and 15%, respectively) and *A. phagocytophilum* (14% and 2%, respectively). *B. miyamotoi* and POWV were detected only in adult ticks and at lower prevalences (3% and 2%, respectively) (Table 1; Fig. 2). We performed additional assays to characterize the strains of *A. phagocytophilum* and POWV. There are two strains of *A. phagocytophilum* that differ by 2 bp from their 16S rRNA sequences. One variant (AP variant 1–nonpathogenic) does not infect humans and is carried by deer (47, 48). The HA strain that is implicated in human disease was present in 67 out of 90 (75%) *A. phagocytophilum*-positive ticks; 19 were found to be of the nonpathogenic strain (21%), and 4 could not be identified further. All POWV strains were part of lineage II, which is associated exclusively with transmission by *I. scapularis*. In addition to these five agents, we found that of the 187 ticks tested, 87% were positive for *Rickettsia*. Of sequences of PCR products from a 20%




FIG 2 Prevalence of five pathogens in *I. scapularis* adults and nymphs.

subset of the *Rickettsia*-positive ticks, all were identified as *R. buchneri*, an ovarian symbiont. Overall, there were no differences in regional infection rates, with the exceptions of *B. microti* ($P = 0.02$) and POWV ($P = 0.01$), which were more prevalent in northern Suffolk County (Table 1; Fig. 2). There were no statistically significant differences in infection rates between spring and fall collections in all the pathogens tested for *I. scapularis* (Table 2). With the exceptions of *R. buchneri* in *I. scapularis* ticks and *R. amblyommatis* in lone star ticks, adult ticks had significantly higher infection rates with *B. burgdorferi* and *A. phagocytophilum* than nymphs ($P = 0.00003$ and $P = 0.004$, respectively) (Tables 1, 2, and 5; Fig. 2). This is a very well-known feature of tick-borne diseases.

A total of 22% (147 of 678) of *I. scapularis* ticks were infected with more than one pathogen. Dual infections were detected in 126 (19%) of the ticks. Triple infections were observed in 21 (3%) of the ticks. A single adult tick collected in the fall was coinfecting with four pathogens (Table 3). The highest coinfection prevalence was found with *B. burgdorferi* and *B. microti*, with 9% of the total ticks analyzed. Ticks coinfecting with *B. burgdorferi* and *A. phagocytophilum* accounted for 7% of the total ticks. Triple infections with these three pathogens were more prevalent in both seasons than dual infections with *B. burgdorferi*-*B. miyamotoi*, *B. burgdorferi*-POWV, *B. miyamotoi*-*A. phagocytophilum*, and *B. miyamotoi*-*B. microti* (Table 3).

Coinfection of *A. phagocytophilum* and *B. microti* with *B. burgdorferi* in host-seeking *I. scapularis* ticks has been reported in several regions of the United States (Table 4). The 54%, 14%, and 13% prevalences of *B. burgdorferi*, *B. microti*, and *A. phagocytophilum*, respectively, in this study are consistent with the prevalence rates found in New York State (39, 49, 50) but are higher than those of other regions in the Northeast (Table 4). The insularity of Long Island could explain and be a determinant of the high global prevalence rate of infected *I. scapularis* ticks found in this study, since rodent population densities inhabiting this region could be affected by the island syndrome (51) that favors increases in rodent populations, which, in turn, could lead to greater opportunities for juvenile stages of *I. scapularis* to find a rodent host and therefore get infected by one or more pathogens.

Pathogen burden of *A. americanum*. The aggressive species *A. americanum* is implicated in the transmission of the causative agents of ehrlichiosis (Table 5). We detected *Ehrlichia* spp. in 23 (7%) of adult *A. americanum* ticks. *E. ewingii* was the predominant species in our ticks (12 ticks; 4%), followed by *E. chaffeensis* (6 ticks; 2%), and an *Ehrlichia ruminantium*-like species (5 ticks; 1%). We also detected *Ehrlichia* in 4 out of 354 individual *A. americanum* nymphs. Two nymphs were positive for *E. ewingii* and 2 for *E. chaffeensis*. All pools of *A. americanum* nymphs were negative for *Ehrlichia*.

B. lonestari was present in 8 ticks (1%). We did not detect *B. burgdorferi* in *A. americanum*, reasserting that this tick species does not play a role in its transmission. Early reports of the Lyme disease agent in this tick may have been mistaken with *B.*

TABLE 2 Prevalence of five pathogens and *R. buchneri* in *I. scapularis* ticks by geographical region of Suffolk County^a

Agent ^b	No. of positive ticks (% [95% CI])			
	Spring		Fall	
	North (n = 125 [80 A/45 N])	South (n = 73 [60 A/13 N])	North (n = 240)	South (n = 240)
<i>Borrelia burgdorferi</i>				
A	58 (72 [61–82])	27 (45 [32–58])	139 (58 [51–64])	127 (53 [46–59])
N	12 (27 [15–42])	4 (31 [10–61])		
<i>Borrelia miyamotoi</i>				
A	3 (4 [1–11])	1 (2 [0.1–10])	9 (4 [2–7])	7 (3 [1–6])
N	0 (0 [0–10])	0 (0 [0–28])		
<i>Anaplasma phagocytophilum</i>				
A	11 (14 [7–24])	8 (13 [6–25])	35 (15 [10–20])	35 (15 [10–20])
N	1 (2 [0.1–13])	0 (0 [0–28])		
<i>Babesia microti</i>				
A	19 (24 [15–35])	8 (13 [6–25])	35 (15 [10–20])	26 (11 [7–16])
N	9 (20 [10–35])	0 (0 [0–28])		
POWV				
A	5 (6 [2–15])	0 (0 [0–7])	5 (2 [1–5])	1 (0.4 [0–3])
N	0 (0 [0–10])	0 (0 [0–28])		
<i>Rickettsia buchneri</i> ^c				
A	18 (78 [56–92])	8 (89 [51–99])	29 (91 [74–97])	35 (87 [72–95])
N	21 (91 [70–98])			

^aValues are means and 95% confidence intervals (CI).

^bA, adults; N, nymphs.

^cThirty-two adults and 23 nymphs were tested for the spring season; 72 adult ticks were tested for the fall season.

lonestari (52). None of the *A. americanum* ticks had detectable *B. microti*, *B. miyamotoi*, *A. phagocytophilum*, or POWV.

We detected a single *Rickettsia* species in *A. americanum*, identified as *R. amblyommatis*. More than half of *A. americanum* (58%) ticks were positive for *R. amblyommatis*.

Impact of tick-borne blood infections. Three pathogens detected in this study infect blood cells. *A. phagocytophilum* infects neutrophils, *B. microti* infects erythrocytes, and the *Ehrlichia* spp. infect monocytes. All three pathogens can also be transmitted by transfusion of blood and blood products (53–55).

TABLE 3 Polymicrobial infections detected in *I. scapularis* ticks

Polymicrobial infections	No. (%) of coinfecting <i>I. scapularis</i> ticks		
	Spring (n = 198)	Fall (n = 480)	Total (n = 678)
Two pathogens			
<i>B. burgdorferi</i> - <i>B. miyamotoi</i>	2 (1)	6 (1)	8 (1)
<i>B. burgdorferi</i> - <i>A. phagocytophilum</i>	11 (6)	39 (8)	50 (7)
<i>B. burgdorferi</i> - <i>B. microti</i>	20 (10)	41 (8)	61 (9)
<i>B. burgdorferi</i> -POWV	2 (1)	1 (0.2)	3 (0.4)
<i>B. miyamotoi</i> - <i>A. phagocytophilum</i>		3 (0.6)	3 (0.4)
<i>B. miyamotoi</i> - <i>B. microti</i>		1 (0.2)	1 (0.1)
Three pathogens			
<i>B. burgdorferi</i> - <i>B. miyamotoi</i> - <i>A. phagocytophilum</i>		2 (0.4)	2 (0.3)
<i>B. burgdorferi</i> - <i>A. phagocytophilum</i> - <i>B. microti</i>	5 (2)	2 (0.4)	7 (1)
<i>B. burgdorferi</i> - <i>B. microti</i> -POWV	2 (1)	1 (0.2)	3 (0.4)
<i>A. phagocytophilum</i> - <i>B. microti</i> -POWV		1 (0.2)	1 (0.1)
Four pathogens			
<i>B. burgdorferi</i> - <i>B. miyamotoi</i> - <i>A. phagocytophilum</i> - <i>B. microti</i>		1 (0.2)	1 (0.1)

TABLE 4 Prevalence of the most common single infections and pathogen coinfections reported in questing *I. scapularis* ticks in the United States from 2003 to 2017

Location	Yr(s)	No. of ticks	Stage ^a	Single infections (%) ^b			Double infections (%)		Triple infections (%) of Bb-Apc-Bm	Reference
				Bb	Ap ^c	Bm	Bb-Ap ^c	Bb-Bm		
NJ	2003–2004	147	A	50	6		3		115	
NY	2003–2006	3,300	N	14	6	3	0.5	1		
		7,914	A	46	12	2	6	1	0.4	
ME	2003	100	A	58	16	7	9	11 ^d	117	
IN	2004	100	A	72	5		4	4 ^d	117	
PA	2005	94	A	52	1		1	1 ^d	117	
WI	2006	100	A	35	14		8	4 ^d	117	
MI	2006	119	A	50	4		2		118	
NJ	2004–2007	478	N	10		4	3			
		610	A	45		8	6			
IA	2007–2009	156	N	17	29		6		120	
NY	2008	132	A	62	22	26	16	22	7	
CT	2008	154	A	65	17	16	16	12	3	
WI	2009–2013	748	N	29	5		3		121	
NY	2011	323 ^e	N	67	34	19				
		922 ^e	A	60	23	4				
		466	A	55	18	3	10	1	1	
NY	2011–2012	4,368	N	19	5	6	2	7	1	
CT	2011–2012	514	N	13	3	6	1	2		
MD	2011–2012	124	N	19	1					
NY	2011–2012	207	N	23	5	11		6		
PA	2013	1,363	A	47	3	3	1	2		
MD	2014–2015	168	N	21	1					
NY	2014–2015	299	N	17	9	3	3	4		
PA	2014–2015	114	N	22	3		2			
VA	2014–2015	472	N	12	1		1			
DC	2014–2015	253	N	23	4					
ME	2015	154	N	18	3	4		3		
MN	2015	1,240	N	25	6	5	2	2	1	
WI	2015	112	A	41	11	9	1	1	1	
PA	2015–2017	1,721	N	25	1	3	1	1		
NY	2016–2017	197	A	56	11	8	4	0.5	2	

^aN, nymphs; A, adults.

^bBb, *Borrelia burgdorferi*; Ap, *Anaplasma phagocytophilum*; Bm, *Babesia microti*.

^cAp human variant.

^dCoinfection of *B. burgdorferi* and *B. microti* plus *Babesia odocoilei* combined.

^eTested in pools of single individuals to a maximum of 10.

The ratio of Lyme disease cases to babesiosis cases has been approximately 4 to 1 for the last 8 years, higher than in other areas where *I. scapularis* is endemic (Fig. 3; Table 4). This ratio also holds for the ratio of *B. burgdorferi* to *B. microti* in ticks in this study, i.e., 3.8 to 1 (Tables 1 and 2). Conversely, during the same 8-year period, the ratio of Lyme disease cases to cases of anaplasmosis is approximately 15 to 1, but our data showed a 4-to-1 ratio of *B. burgdorferi* to *A. phagocytophilum*. This disparity between the reported cases of anaplasmosis and the tick infection rates is not readily explainable but could be due to the role that the nonpathogenic strain might have in exposure. Alternatively, a larger number of anaplasmosis cases may not be severe enough to warrant medical attention.

There are increasing numbers of reported cases of ehrlichiosis in Suffolk County (Fig. 3). Although the infection rate of *Ehrlichia* spp. in *A. americanum* nymphs and adults was less than 5% (Table 5), the growing populations of these ticks as well as their aggressive nature create additional anxiety over this emerging infection (Fig. 3).

The three tick-borne blood infections have several features in common. The frequency of reported cases of all three is highest in the elderly (anaplasmosis susceptibility appears to be less bound by age) and in males. Importantly, people at risk for clinical infection and for increased disease severity have compromised immune systems due to a number of causes, which can include cancer therapy, immunosuppressive

TABLE 5 Agents detected in *A. americanum* and *D. variabilis* ticks^a

Tick species and agent ^b	No. of positive ticks (% [95% CI])		
	Spring		Total
	North	South	
<i>Amblyomma americanum</i>	<i>n</i> = 352 (128 A/224 N)	<i>n</i> = 324 (194 A/130 N)	<i>n</i> = 676 (322 A/354 N)
<i>Borrelia lonestari</i> ^c			
A	1 (1 [0.05–6])	5 (3 [1–8])	6 (2 [1–5])
N	1 (0.5 [0.2–3])	1 (1 [0.5–6])	2 (0.6 [0.1–3])
<i>Ehrlichia chaffeensis</i>			
A	3 (2 [0.6–7])	3 (1 [0.4–5])	6 (2 [0.7–4])
N	1 (0.4 [0.2–3])	1 (0.8 [0.4–5])	2 (0.6 [0.1–2])
<i>Ehrlichia ewingii</i>			
A	8 (6 [3–12])	4 (2 [0.7–5])	12 (4 [2–6])
N	2 (1 [0.1–3])	0 (0 [0–3])	2 (0.6 [0.1–2])
<i>E. ruminantium</i> -like species			
A	2 (2 [0.3–6])	3 (1 [0.4–5])	5 (1 [0.6–4])
N	0 (0 [0–2])	0 (0 [0–3])	0 (0 [0–1])
Total <i>Ehrlichia</i> spp.			
A	13 (10 [6–17])	10 (5 [3–9])	23 (7 [5–11])
N	3 (1 [0.3–4])	1 (0.8 [0.4–5])	4 (1 [0.4–3])
<i>Rickettsia amblyommatis</i>			
A	79 (62 [53–70])	119 (61 [54–68])	198 (61 [56–67])
N	111 (49 [43–56])	85 (65 [56–73])	196 (55 [50–60])
<i>Dermacentor variabilis</i>	<i>n</i> = 124	<i>n</i> = 172	<i>n</i> = 296
<i>Rickettsia montanensis</i>	6 (5 [2–11])	2 (1 [0.2–5])	8 (3 [1–5])

^aValues are means and 95% confidence intervals (CIs).

^bA, adult; N, nymph.

^cA total of 548 ticks were tested from the northern (102 adults, 210 nymphs) and southern (140 adults, 96 nymphs) regions of Suffolk County.

drugs to treat autoimmune disorders, splenectomy, preexisting liver disease, and endogenous disorders of the immune system, among others. In communities with both high tick infestations and the presence of a sizable elderly population, the features of the three blood infections are important contributors to their high morbidity and mortality (<https://www.cdc.gov/anaplasmosis/stats/index.html>; <https://www.cdc.gov/parasites/babesiosis/disease.html>; <https://www.cdc.gov/ehrlichiosis/stats/index.html>).

The genus *Rickettsia* constitutes the largest biomass in the three tick species.

R. montanensis (2.7%) was the only agent detected in *D. variabilis* (Table 5). The cases of RMSF have decreased markedly in Suffolk County since the 1970s, with the numbers of cases ranging from 2 to 14 per year in the last decade (<https://www.health.ny.gov/statistics/diseases/communicable/>). *R. amblyommatis* was present in more than half of the *A. americanum* ticks sampled in this study (Table 5). *R. buchneri* is an ovarian endosymbiont with an overwhelming presence in *I. scapularis* (Table 1; Fig. 2) and has not been associated with human disease. However, its presence could be a factor in the associations with pathogens found simultaneously in this tick.

DISCUSSION

The polymicrobial infection approach to tick-borne diseases is rooted in the discovery of new or emergent pathogens, such as the ones detected in this study, changes in or discovery of new pathogen genotypes, such as is the case for POWV and *A. phagocytophilum*, and expansion of tick ranges as well as of their pathogens, as represented by *A. americanum*. The changes in tick populations and their pathogens may be the result of climate change, with unpredictable short- or long-term consequences. However, factors other than climate change, such as anthropogenic changes and host availability, might have great impact.

***Ixodes scapularis* microbiome.** *I. scapularis* is a public health threat (56). *B. burgdorferi* infects more than half of the adult *I. scapularis* ticks tested, and this accounts for the high number of Lyme disease cases in our area (Fig. 3). Of note is that nearly one-quarter of the *I. scapularis* ticks tested had polymicrobial infections, and this justifies the modification of the clinical approach to tick-borne diseases to cover all




FIG 3 Number of cases of anaplasmosis, babesiosis, ehrlichiosis, and Lyme disease from 2010 to 2017 in Suffolk County, New York. Numbers were derived from <https://www.health.ny.gov/statistics/diseases/communicable/>.

infection possibilities. At a more fundamental level, the polymicrobial infections open new possibilities for research into the microbial relationships with and within the tick.

Babesia microti causes clinical disease in elderly, immunosuppressed, and splenectomized patients (57, 58) and also in blood transfusion recipients (59). Asymptomatic babesiosis exists in individuals that do not have the above risk factors, representing a group of seropositives, and these represent a major risk as blood transfusion donors. Genomic tools have added new perspectives on the expansion of *B. microti* (60). There are Northeast and Midwest lineages segregating into local populations of the parasite (61, 62), suggesting dynamic adaptation changes promoting expansion.

Anaplasmosis is variable in terms of severity. The two variants of *A. phagocytophilum* also have ecological implications, in that the AP-HA variant is acquired from mice, whereas the reservoir of the nonpathogenic variant is deer. That these variants were detected in our study is representative of the advances at the molecular biology level that have a bearing on epidemiology.

POWV is the causative agent of a life-threatening encephalitis (63). There have not been any reported cases of POWV in our area, but its low-level presence (lineage II) in *I. scapularis* could result in transmission to humans. Lineage II virus circulates in an *I. scapularis*–white-footed mouse cycle, and it is precisely in this cycle that this virus can become a part of the polymicrobial group of pathogens (63–69).

Amblyomma americanum expansion. Since the first confirmed records (22), and thereafter (23, 70), the presence of *A. americanum* has increased markedly, and it is likely the most abundant tick species in our environment. However, the lack of systematic tick surveillance in the intervening years since the first records were made makes this difficult to confirm. The expansion of *A. americanum* into the North Atlantic states has been mapped and documented (20, 21, 52, 71). Sonenshine connected the northward expansion of *A. americanum* to climate change (72). This expansion has occurred in a short time. An increase of nearly 60% in *A. americanum* numbers has been reported in Connecticut in the last 20 years (73), as well as in New Jersey (74). Studies

in our laboratory showed that genetic changes occur in the tick population during range expansion (75).

There are serious consequences to the range expansion of *A. americanum*. Notably, even in the absence of transmitted pathogens, this species is an aggressive and tenacious tick. All three stages bite humans, and the larvae, in particular, can infest in large numbers, leading to an uncomfortable dermatitis. There are some subtle effects as well. Large populations of *A. americanum* in a community can lead to an exaggerated perception of the risk of Lyme disease, as most people cannot differentiate between *I. scapularis* and lone star ticks (76).

The cases of ehrlichiosis have increased steadily in our area in recent years as a function of the increases in the populations of *A. americanum* (Fig. 3). Our results indicate that there are three *Ehrlichia* species in Suffolk County. *E. ewingii* is the most frequent *Ehrlichia* species present in our ticks, and therefore, it could be the main causative agent of human ehrlichiosis in this area (49) (Table 5). *E. chaffeensis* is clinically indistinguishable from *E. ewingii*, and species identification is not possible by serology, so it may be difficult to determine the infecting species (77). An *E. ruminantium*-like species is present in *A. americanum* in our area, although its disease-causing potential is unclear (78). Transfusion-acquired *E. ewingii* has been documented (79), making this organism a serious threat to recipients who are at greater risk of a severe infection (80), and infection with *E. ewingii* has been documented in patients that were not immunosuppressed. Pediatric ehrlichiosis with an increased case fatality rate has been documented as well (81).

B. lonestari has been detected at low levels, but STARI has not been documented in our area as it has elsewhere in areas where *A. americanum* is abundant (82–84). *B. lonestari* has been linked as the causative agent of STARI from a single patient with erythema migrans (85). However, to date, this has been the only case reported.

Rickettsial biome. The rickettsiae constitute the largest prokaryotic biomass in two of the three species of ticks detected here. The sheer abundance of these rickettsiae can be expected to have a major role in the vector-pathogen relationship and to influence the pathogen-patient relationship after transmission.

There was overwhelming infection of *R. amblyommatis* (40) in our *A. americanum* populations. This rickettsia may be more than a symbiont, as cases of rickettsiosis may have been caused by *R. amblyommatis* (86). RMSF-like illness may be associated with the expanding range of *A. americanum* (21). This is further supported by the lack of detection of *R. rickettsii* in tick surveillance studies, including the work presented here. However, that there is a high prevalence of *R. amblyommatis* does not necessarily support its role in human disease. Whether a symbiont, a pathogen under certain conditions, or an outright pathogen, *R. amblyommatis* has introduced a confounding factor in the diagnoses of febrile tick-borne infections, as positive serologies may be misinterpreted.

The rickettsiae present in the ovaries of *I. scapularis* belong to the spotted fever group and are closely related to *R. montanensis* (36, 87, 88). The most abundant prokaryotic DNA sequences found in *I. scapularis* were from *R. buchneri* (37–39). *R. buchneri* underwent the introduction of transposons with genomic deletions and mutations that resulted in the loss of pathogenicity (89). *R. buchneri* contains multiple mobile genetic elements that endow this species with a plastic and repetitive genome that is thought to account for its symbiotic lifestyle (90). Its overwhelming presence in this study suggests that this organism plays an undisputed role in the homeostasis of this tick.

R. montanensis was the only agent detected in *D. variabilis* ticks, with no detection of *R. rickettsii*. This tick was associated with an outbreak of RMSF in our area in the late 1970s (91). Given our findings in this study, we support the suggestion that other rickettsial species could confound a diagnosis of RMSF (92, 93), although the prevalence of this symbiotic organism in *D. variabilis* seems to have remained stable for many years (92).

Polymicrobial infections may be synergistic in enhancing the severity of human illnesses, and more-deleterious interactions could be discovered if there was greater emphasis on a pluralistic approach to tick-borne diseases. There is an extensive literature on the topic of coinfections, and we cite the most recent for American patients (94) and for Europe (95). However, despite the high levels of tick coinfections, there is at least one therapeutic feature that could mitigate the impact of polymicrobial infections in patients. Two of the most common coinfecting organisms, *B. burgdorferi* and *A. phagocytophilum*, are responsive to doxycycline, so both could be treated simultaneously. Polymicrobial infections can lead to treatment problems as well. The possibility of side effects has resulted in doxycycline not being prescribed to young children, an overrepresented group in Lyme disease. *B. microti* requires antiparasitic drugs that do not work against *B. burgdorferi* and *A. phagocytophilum*. Likewise, beta-lactam antibiotics can be active against *B. burgdorferi* but not *A. phagocytophilum*, and it is in these cases where the complexity of polymicrobial infections requires further treatment evaluation.

Nonetheless, the relationships of pathogens and symbionts in the tick have biological significance but appear to be complex. Pathogen burden could be important in maintaining the balance of organisms within the tick, where deviations may alter the mechanics of transmission. On one hand, there is evidence for the possibility of limited interactions among the several coinfecting pathogens in *I. scapularis*. These pathogens occupy different anatomical niches in the tick, as some are extracellular in the lumen of the midgut (*Borrelia*) and others are obligate intracellular organisms (*Anaplasma*, *Babesia*, and POWV), so it is possible that direct contact among them may be limited. There is a hierarchy of transmission of pathogens in relation to the duration of blood feeding. POWV can be transmitted within 15 min of tick attachment, and both *A. phagocytophilum* and *B. miyamotoi* can be transmitted within the first 24 h of attachment. Transmission of *B. burgdorferi* increases with the length of attachment (96–98), and *B. microti* needs to change its morphology while in the salivary glands of the tick, requiring a longer period of time for transmission. This pattern of staggered transmission may further segregate the pathogens within the vector and actually favor acquisition of the organisms that are transmitted quickly, as people may remove ticks in time to abort infections with the slower pathogens.

Notwithstanding, there is increasing evidence for far-reaching interactions among symbionts and pathogens with each other and, collectively, with the ticks. In some instances, the interaction can be to the benefit of the pathogen. *A. phagocytophilum* induces the production of the antifreeze glycoprotein (IAFGP), which in turn makes infection easier (99). Relationships between the symbiont-pathogen and the tick may result in a neutral status quo; for example, the vector competence of *A. americanum* for *R. rickettsii* was not significantly affected by *R. amblyommatidis* (100). In other instances, tick responses can be harmful to pathogens. Hemocytes of *I. scapularis* ingest *B. burgdorferi* (101), and this tick regulates infection with *A. phagocytophilum* through the production of antimicrobial peptides, and actin phosphorylation for survival, as well as proteases to increase vector fitness (102–104). *R. montanensis* (an organism of unproven pathogen status) induces ticks to produce defensins (105) and protease inhibitors that limit colonization by this *Rickettsia* species (106). *Anaplasma*, *Borrelia*, *Ehrlichia*, and *Rickettsia* spp. do not have interbacterial effector and immunity genes whose products regulate interactions among bacteria utilizing the same niche (38). The lack of these genes can lead to a system of shared tolerance for each other that would be clearly to their advantage for survival in the vector. It appears that we have only begun to appreciate the interactions of the microbes with each other and with the tick. The best approach to study the interactions among the prokaryotes in the tick is to consider the polymicrobial nature of these fascinating biological systems (107–109).

MATERIALS AND METHODS

Tick collection and study areas. Two active tick-borne pathogen surveillance programs were conducted during the spring season peak (from May to July) and the fall season peak (from October to

TABLE 6 Primer and probe sequences for RT-PCR^a

Pathogen	Gene target	Primer pair	Probe	5' dye	3' quencher
<i>Borrelia burgdorferi</i>	<i>ospA</i>	Fwd: CCTTCAAGTACTCCAGATCCATTG Rev: AACAAAGACGGCAAGTACGATC	CAACAGTAGACAAGCTTGA	6-FAM	MGB
<i>Borrelia miyamotoi</i> / <i>B. lonestari</i>	<i>flaB</i>	Fwd: AGCACAAAGCTTCATGGACATTGA Rev: GAGCTGCTTGAGCACCTTCTC	TGTGGGTGCAAATCAGGATGAAGCA	HEX	BHQ-1
<i>Babesia microti</i>	<i>cox1</i>	Fwd: CATCATGCCAGGCCTGTTTG Rev: GAAGAAACCACAAGAGCAAATGC	TACTACCCATACTGGTCGGTGCTCC	Quasar 705	BHQ-2
<i>Anaplasma phagocytophilum</i>	16S rRNA	Fwd: GGCATGTAGGCGGTTCCGGT Rev: CACTAGGAATCCGCTATCCTCTCC	GCCAGGGCTTAACCCTGGAGCT	Cy5	BHQ-2
Powassan virus	3'UTR	Fwd: GTGATGTGGCAGCGCACC Rev: CTGCGTCGGGAGCGACCA	CCTACTGCGGCAGCACACACAGTG	Texas Red	BHQ-2
<i>Ehrlichia</i> spp.	16S rRNA	Fwd: CGTAAAGGGCAGCTAGGTGGACTA Rev: CACCTCAGTGTCAATCGAACCA	TCGAAAGAGGATAGCGGA	VIC	MGB
<i>Rickettsia</i> spp.	<i>ompB</i>	Fwd: AACAAAGCTGCTGGGCACCATAT Rev: CGGTCTGCTATCGGTATCACT	AGAGAATGAGAAACCGTTAACGT	FAM	MGB

^aUTR, untranslated region; 6FAM, 6-carboxyfluorescein; HEX, 6-carboxy-2,4,4,5,7,7-hexachlorofluorescein; BHQ, black hole quencher.

November) throughout Suffolk County, New York, in 2018. Questing ticks were collected from vegetation along trails by flagging a 1-m² cotton flannel fabric attached on both ends to a wooden pole between 10:00 and 14:00 h during sunny days. The area of study was divided into two geographical regions to compare potential differences in infection rates between the northern and the southern regions of the county (Fig. 1). Ticks were collected for a minimum of 60 min per site in order to collect a representative number of each species to estimate the prevalence of pathogens in each geographical region.

All collected ticks were identified morphologically to species, life stage, and sex by use of a dissecting microscope and the appropriate taxonomic keys and stored at -80°C until further processing (110–112).

TNA extraction from ticks. Total nucleic acids (TNA) were extracted using the NucliSENS easyMAG platform (bioMérieux, Durham, NC). Ticks were grouped by location and day of collection. To remove environmental contaminants, pools of 10 ticks were washed with 3% hydrogen peroxide, followed by three washes with 1 ml of $1\times$ phosphate-buffered saline (PBS). The ticks were then placed individually in a sterile 1.5-ml centrifuge tube and homogenized using a 21-gauge, 1.5-in needle in 50 μl of $1\times$ PBS. The entire volume was then added to NucliSENS easyMAG lysis buffer (bioMérieux, Durham, NC). TNA were extracted according to the manufacturer's protocol, eluted in 40 μl , and stored at -80°C .

Pathogen detection by real-time PCR. We utilized a pathogen detection strategy in which we tested each tick TNA by quantitative PCR (qPCR) using two approaches, consisting of either a single-agent qPCR using DNA as a template or a multiplex one-step reverse transcription PCR (RT-qPCR), using both cDNA and DNA as a template (113).

All *I. scapularis* and *A. americanum* samples were screened for the presence of *B. burgdorferi*, *B. miyamotoi*/*B. lonestari*, *B. microti*, *A. phagocytophilum*, and POWV with the multiplex RT-qPCR (Table 6). Our rationale for testing *A. americanum* was to evaluate the role of this tick as a potential vector of these five agents. For positive controls of the multiplex assay, we employed quantified plasmid standards at 10, 100, and 1,000 copies (113). All remaining tests consisted of single-agent qPCR assays. For detection of *Rickettsia*, we developed a qPCR assay designed to detect a fragment within the *ompB* gene of the most common spotted fever group *Rickettsia* species. This assay was used to screen all TNA from *A. americanum* and *D. variabilis* ticks and from 127 *I. scapularis* ticks. For positive controls, we used TNA from an *I. scapularis* adult previously shown to be infected with *R. buchneri*. To detect *Ehrlichia* spp., we employed an *Ehrlichia*-specific assay targeting a portion of the 16S rRNA gene. This assay was used to test all *A. americanum* ticks (Table 6). For positive controls, we used DNA from a lysate of *E. chaffeensis*-infected DH82 cells.

All qPCRs were performed on a Bio-Rad C1000 Touch system with a CFX96 optical module (Bio-Rad, Hercules, CA) using the RNA UltraSense one-step quantitative RT-PCR system (Invitrogen, Carlsbad, CA). The final reaction mixture contained 5 μl of template and 20 μl of master mixture. The master mixture contained 0.2 μM each forward primer, 0.3 μM each reverse primer, 0.1 μM each probe, 1.25 μl of RNA UltraSense enzyme mix, and 5 μl RNA UltraSense $5\times$ reaction mix. The reverse transcription step was performed at 55°C for 15 min, followed by incubation at 95°C for 10 min. The RT step was omitted for the *Ehrlichia* and *Rickettsia* qPCR assays. The PCR consisted of 40 cycles at 95°C for 15 s and 60°C for 30 s.

Qualitative PCR. We developed qualitative PCR assays to further characterize the species detected by qPCR. For *Rickettsia*, we employed an assay that amplified 380 nucleotides (nt) within *ompB*. For *Anaplasma*, we designed an assay within *gltA* to differentiate between the pathogenic agent (AP-HA) and nonpathogenic variants (AP-variant 1). For *Ehrlichia*, we first employed an assay targeting the 16S rRNA gene. Any samples found to be positive for the *E. ruminantium* variant were tested with a nested PCR targeting the *gltA* gene.

TABLE 7 Primer sequences and PCR conditions for qualitative assay

Pathogen	Gene target	Primer pair	Product length (nt)	PCR conditions
<i>Borrelia burgdorferi</i>	<i>ospA</i>	Fwd: GCGTTTCAGTAGATTTGCCT Rev: TTGGTGCCATTTGAGTCGTA	676	95°C for 10 min; 40 cycles of 95°C for 30 s, 57°C for 30 s, and 72°C for 60 s
<i>Borrelia miyamotoi</i> / <i>B. lonestari</i>	<i>flaB</i>	Fwd: GGGATTATMAATCATAATACRTCAGC Rev: TTGCTTGTGCAATCATAGCCATTGC	967 (<i>B. miyamotoi</i>)/ 949 (<i>B. lonestari</i>)	95°C for 10 min; 40 cycles of 95°C for 30 s, 57°C for 30 s, and 72°C for 70 s
<i>Babesia microti</i>	18S rRNA	Fwd: GGGACTTTGCGTTCATAAAACGC Rev: GCAATAATCTATCCCCATCACGAT	171	95°C for 10 min; 40 cycles of 95°C for 30 s, 60°C for 30 s, and 72°C for 45 s
<i>Anaplasma phagocytophilum</i>	<i>gltA</i>	Primary reaction Fwd: ACCGGAACCCCATAGCTCT Rev: GCAAGTCGATTGATCCGCT	668	95°C for 10 min; 40 cycles of 95°C for 30 s, 62°C for 30 s, and 72°C for 60 s (same conditions for primary and secondary reactions)
		Secondary reaction Fwd: TCGACATTTGGGTACAACCTTGCG Rev: CAATTGCGAAAGTACCCGGCA	589	
Powassan virus	Envelope (E) glycoprotein	Fwd: GGCAACTGCATCTCTATRAATCC Rev: CCTCATGCAGTGAAAATGGATATCTT	395	95°C for 10 min; 40 cycles of 95°C for 30 s, 58°C for 30 s, and 72°C for 45 s
<i>Ehrlichia</i> spp.	16S rRNA	Fwd: ATGCGTAGGAATCTACCTAGTAGTA Rev: GCCTTGGTATTTCACTTTAACTTACT	460	
<i>Ehrlichia</i> spp.	<i>gltA</i>	Primary reaction Fwd: CCAGGATTTATGTCTACTGCTGC Rev: GCATACYCTATGACCAAMCCCAT	777	95°C for 10 min; 40 cycles of 95°C for 30 s, 56°C (primary)/58°C (secondary) for 30 s, and 72°C for 60 s
		Secondary reaction Fwd: GCGCGGATTACRTTTATTGATGG Rev: ATTGGCHCCACCATGAGCTG	627	
<i>Rickettsia</i> spp.	<i>ompB</i>	Fwd: GGTACTGCCGAGTTACGTTTAG Rev: CTCGCATCAACAACRCCTG	380	95°C for 10 min; 40 cycles of 95°C for 30 s, 57°C for 30 s, and 72°C for 45 s

For characterization of *B. burgdorferi*, *B. miyamotoi*, *B. lonestari*, and POWV, we used our previously established assays (49, 114).

All qualitative PCRs were performed using AmpliTaq Gold 360 master mix (Applied Biosystems, Foster City, CA) in a 25 μ l-reaction mixture with 2 μ l of template, 0.2 μ M forward primer, and 0.3 μ M reverse primer and 12.5 μ l of the polymerase. Primers and reaction conditions used for each pathogen are described in Table 7. A nontemplate control was included in each assay. PCR products were resolved through 2% agarose gel in Tris-borate-EDTA (TBE) buffer and sequenced by dideoxy sequencing. All sequences were analyzed using Geneious 10.0.9 software.

Statistical analysis. A Fisher's exact test was performed for each agent to compare seasonal and geographical infection rates with the five pathogens. A two-tailed *P* value of <0.05 was considered statistically significant. Statistical analyses were performed using the software package R (R-Development Core Team; www.r-project.org). The mean and 95% confidence intervals of pathogen prevalence were calculated using the proportion test in R.

ACKNOWLEDGMENTS

DNA from cell-cultured *E. chaffeensis* was used as a positive control for PCR and was provided by J. Stephen Dumler at the Walter Reed National Military Medical Center. We thank the New York State Office of Parks, Recreation and Historic Preservation and the U.S. National Park Service for permission to collect ticks. The assistance of Erik Kopping, Peter T. Benziger, and Indra Jayatilaka is gratefully acknowledged.

This study was funded by a grant from the National Institutes of Health (AI-027044-29 to J.L.B.). Support was also provided by the Island Outreach Foundation, Blue Point, NY, to the Stony Brook Renaissance School of Medicine. Support from the Steven & Alexandra Cohen Foundation (CU18-2692) was provided to R.T.

REFERENCES

- Telford SRT, Goethert HK. 2008. Emerging and emergent tick-borne infections, p 344–376. In Bowman AS, Nuttall PA (ed), *Ticks: biology, disease and control*. Cambridge University Press, Cambridge, United Kingdom.
- Rosenberg R, Lindsey NP, Fischer M, Gregory CJ, Hinckley AF, Mead PS, Paz-Bailey G, Waterman SH, Drexler NA, Kersh GJ, Hooks H, Partridge SK, Visser SN, Beard CB, Petersen LR. 2018. Vital signs: trends in reported vectorborne disease cases—United States and Territories, 2004–2016. *MMWR Morb Mortal Wkly Rep* 67:496–501. <https://doi.org/10.15585/mmwr.mm6717e1>.
- Fukunaga M, Takahashi Y, Tsuruta Y, Matsushita O, Ralph D, McClelland M, Nakao M. 1995. Genetic and phenotypic analysis of *Borrelia miyamotoi*.

- amotai sp. nov., isolated from the ixodid tick *Ixodes persulcatus*, the vector for Lyme disease in Japan. *Int J Syst Bacteriol* 45:804–810. <https://doi.org/10.1099/00207713-45-4-804>.
4. Telford SR, Armstrong PM, Katavolos P, Foppa I, Garcia AS, Wilson ML, Spielman A, Spielman A. 1997. A new tick-borne encephalitis-like virus infecting New England deer ticks, *Ixodes dammini*. *Emerg Infect Dis* 3:165–170. <https://doi.org/10.3201/eid0302.970209>.
 5. Burgdorfer W, Barbour AG, Hayes SF, Benach JL, Grunwaldt E, Davis JP. 1982. Lyme disease—a tick-borne spirochetosis? *Science* 216:1317–1319. <https://doi.org/10.1126/science.7043737>.
 6. Grunwaldt E. 1977. Babesiosis on Shelter Island. *N Y State J Med* 77:1320–1321.
 7. Ruebush TK, Juranek DD, Chisholm ES, Snow PC, Healy GR, Sulzer AJ. 1977. Human babesiosis on Nantucket Island. *N Engl J Med* 297:825–827. <https://doi.org/10.1056/NEJM197710132971511>.
 8. Filstein MR, Benach JL, White DJ, Brody BA, Goldman WD, Bakal CW, Schwartz RS. 1980. Serosurvey for human babesiosis in New York. *J Infect Dis* 141:518–521. <https://doi.org/10.1093/infdis/141.4.518>.
 9. Benach JL, Habicht GS. 1981. Clinical characteristics of human babesiosis. *J Infect Dis* 144:481. <https://doi.org/10.1093/infdis/144.5.481>.
 10. Swanson SJ, Neitzel D, Reed KD, Belongia EA. 2006. Coinfections acquired from ixodes ticks. *Clin Microbiol Rev* 19:708–727. <https://doi.org/10.1128/CMR.00011-06>.
 11. Chen SM, Dumler JS, Bakken JS, Walker DH. 1994. Identification of a granulocytotropic Ehrlichia species as the etiologic agent of human disease. *J Clin Microbiol* 32:589–595.
 12. Bakken JS, Dumler JS, Chen S-M, Eckman MR, Van Etta LL, Walker DH. 1994. Human granulocytic ehrlichiosis in the Upper Midwest United States. *JAMA* 272:212. <https://doi.org/10.1001/jama.1994.03520030054028>.
 13. Moss WJ, Dumler JS. 2003. Simultaneous infection with *Borrelia burgdorferi* and human granulocytic ehrlichiosis. *Pediatr Infect Dis J* 22:91–92. <https://doi.org/10.1097/0006454-200301000-00023>.
 14. Agüero-Rosenfeld ME, Donnarumma L, Zentmaier L, Jacob J, Frey M, Noto R, Carbonaro CA, Wormser GP. 2002. Seroprevalence of antibodies that react with *Anaplasma phagocytophila*, the agent of human granulocytic ehrlichiosis, in different populations in Westchester County, New York. *J Clin Microbiol* 40:2612–2615. <https://doi.org/10.1128/jcm.40.7.2612-2615.2002>.
 15. Steere AC, McHugh G, Suarez C, Hoitt J, Damle N, Sikand VK. 2003. Prospective study of coinfection in patients with erythema migrans. *Clin Infect Dis* 36:1078–1081. <https://doi.org/10.1086/368187>.
 16. Dumler JS, Barbet AF, Bekker C, Dasch GA, Palmer GH, Ray SC, Rikihisa Y, Rurangirwa FR. 2001. Reorganization of genera in the families Rickettsiaceae and Anaplasmataceae in the order Rickettsiales: unification of some species of Ehrlichia with *Anaplasma*, *Cowdria* with Ehrlichia and Ehrlichia with Neorickettsia, descriptions of six new species combinations and designation of Ehrlichia equi and “HGE agent” as subjective synonyms of Ehrlichia phagocytophila. *Int J Syst Evol Microbiol* 51:2145–2165. <https://doi.org/10.1099/00207713-51-6-2145>.
 17. McLean DM, Donohue WL. 1959. Powassan virus: isolation of virus from a fatal case of encephalitis. *Can Med Assoc J* 80:708–711.
 18. Piantadosi A, Rubin DB, McQuillen DP, Hsu L, Lederer PA, Ashbaugh CD, Duffalo C, Duncan R, Thon J, Bhattacharyya S, Basgoz N, Feske SK, Lyons JL. 2016. Emerging cases of Powassan virus encephalitis in New England: clinical presentation, imaging, and review of the literature. *Clin Infect Dis* 62:707–713. <https://doi.org/10.1093/cid/civ1005>.
 19. Telford SR, Dawson JE, Katavolos P, Warner CK, Kolbert CP, Persing DH, Persing DH. 1996. Perpetuation of the agent of human granulocytic ehrlichiosis in a deer tick-rodent cycle. *Proc Natl Acad Sci U S A* 93:6209–6214. <https://doi.org/10.1073/pnas.93.12.6209>.
 20. Springer YP, Eisen L, Beati L, James AM, Eisen RJ. 2014. Spatial distribution of counties in the continental United States with records of occurrence of *Amblyomma americanum* (Ixodida: Ixodidae). *J Med Entomol* 51:342–351. <https://doi.org/10.1603/me13115>.
 21. Dahlgren FS, Paddock CD, Springer YP, Eisen RJ, Behravesh CB. 2016. Expanding range of *Amblyomma americanum* and simultaneous changes in the epidemiology of spotted fever group rickettsiosis in the United States. *Am J Trop Med Hyg* 94:35–42. <https://doi.org/10.4269/ajtmh.15-0580>.
 22. Means RG, White DJ. 1997. New distribution records of *Amblyomma americanum* (L.) (Acari: Ixodidae) in New York State. *J Vector Ecol* 22:133–145.
 23. Mixson TR, Ginsberg HS, Campbell SR, Sumner JW, Paddock CD. 2004. Detection of Ehrlichia chaffeensis in adult and nymphal *Amblyomma americanum* (Acari: Ixodidae) ticks from Long Island, New York. *J Med Entomol* 41:1104–1110. <https://doi.org/10.1603/0022-2585-41.6.1104>.
 24. Maupin GO, Anderson BE, Piesman JF, Johnson BJB, Sims KG, Happ CM, Olson JG, Childs JE. 1993. *Amblyomma americanum*: a potential vector of human ehrlichiosis. *Am J Trop Med Hyg* 49:239–244. <https://doi.org/10.4269/ajtmh.1993.49.239>.
 25. Buller RS, Arens M, Hmiel SP, Paddock CD, Sumner JW, Rikihisa Y, Unver A, Gaudreault-Keener M, Manian FA, Liddell AM, Schmulewitz N, Storch GA. 1999. *Ehrlichia ewingii*, a newly recognized agent of human ehrlichiosis. *N Engl J Med* 341:148–155. <https://doi.org/10.1056/NEJM199907153410303>.
 26. Ewing SA, Roberson WR, Buckner RG, Hayat CS. 1971. A new strain of Ehrlichia canis. *J Am Vet Med Assoc* 159:1771–1774.
 27. Anderson BE, Greene CE, Jones DC, Dawson JE. 1992. Ehrlichia ewingii sp. nov., the Etiologic agent of canine granulocytic ehrlichiosis. *Int J Syst Bacteriol* 42:299–302. <https://doi.org/10.1099/00207713-42-2-299>.
 28. Walker DH, Ismail N, Olano JP, McBride JW, Yu X-J, Feng H-M. 2004. Ehrlichia chaffeensis: a prevalent, life-threatening, emerging pathogen. *Trans Am Clin Climatol Assoc* 115:375–384.
 29. Barbour AG, Maupin GO, Teltow GJ, Carter CJ, Piesman J. 1996. Identification of an uncultivable *Borrelia* species in the hard tick *Amblyomma americanum*: possible agent of a Lyme disease-like illness. *J Infect Dis* 173:403–409. <https://doi.org/10.1093/infdis/173.2.403>.
 30. Commins SP, James HR, Kelly LA, Pochan SL, Workman LJ, Perzanowski MS, Kocan KM, Fahy JV, Nganga LW, Ronmark E, Cooper PJ, Platts-Mills T. 2011. The relevance of tick bites to the production of IgE antibodies to the mammalian oligosaccharide galactose- α -1,3-galactose. *J Allergy Clin Immunol* 127:1286–93.e6. <https://doi.org/10.1016/j.jaci.2011.02.019>.
 31. Rainey T, Occi JL, Robbins RG, Egizi A. 2018. Discovery of *Haemaphysalis longicornis* (Ixodida: Ixodidae) parasitizing a sheep in New Jersey, United States. *J Med Entomol* 55:757–759. <https://doi.org/10.1093/jme/tjy006>.
 32. Uchida T, Yan Y, Kitaoka S. 1995. Detection of *Rickettsia japonica* in *Haemaphysalis longicornis* ticks by restriction fragment length polymorphism of PCR product. *J Clin Microbiol* 33:824–828.
 33. Luo L-M, Zhao L, Wen H-L, Zhang Z-T, Liu J-W, Fang L-Z, Xue Z-F, Ma D-Q, Zhang X-S, Ding S-J, Lei X-Y, Yu X. 2015. *Haemaphysalis longicornis* ticks as reservoir and vector of severe fever with thrombocytopenia syndrome virus in China. *Emerg Infect Dis* 21:1770–1776. <https://doi.org/10.3201/eid2110.150126>.
 34. Hecht JA, Allerdice MEJ, Dykstra EA, Mastel L, Eisen RJ, Johnson TL, Gaff HD, Varela-Stokes AS, Goddard J, Pagac BB, Paddock CD, Karpathy SE. 3 April 2019. Multistate survey of American dog ticks (*Dermacentor variabilis*) for rickettsia species. *Vector-Borne Zoonotic Dis* <https://doi.org/10.1089/vbz.2018.2415>.
 35. Trout Fryxell RT, Hendricks BM, Pompo K, Mays SE, Paulsen DJ, Operario DJ, Houston AE. 2017. Investigating the adult ixodid tick populations and their associated *Anaplasma*, Ehrlichia, and Rickettsia bacteria at a Rocky Mountain spotted fever hotspot in Western Tennessee. *Vector-Borne Zoonotic Dis* 17:527–538. <https://doi.org/10.1089/vbz.2016.2091>.
 36. Kurtti TJ, Felsheim RF, Burkhardt NY, Oliver JD, Heu CC, Munderloh UG. 2015. *Rickettsia buchneri* sp. nov., a rickettsial endosymbiont of the blacklegged tick *Ixodes scapularis*. *Int J Syst Evol Microbiol* 65:965–970. <https://doi.org/10.1099/ijs.0.000047>.
 37. Moreno CX, Moy F, Daniels TJ, Godfrey HP, Cabello FC. 2006. Molecular analysis of microbial communities identified in different developmental stages of *Ixodes scapularis* ticks from Westchester and Dutchess Counties, New York. *Environ Microbiol* 8:761–772. <https://doi.org/10.1111/j.1462-2920.2005.00955.x>.
 38. Ross BD, Hayes B, Radey MC, Lee X, Josek T, Bjork J, Neitzel D, Paskewitz S, Chou S, Mougous JD. 2018. *Ixodes scapularis* does not harbor a stable midgut microbiome. *ISME J* 12:2596–2607. <https://doi.org/10.1038/s41396-018-0161-6>.
 39. Tokarz R, Tagliaferro T, Sameroff S, Cucura DM, Oleynik A, Che X, Jain K, Lipkin WI. 2019. Microbiome analysis of *Ixodes scapularis* ticks from New York and Connecticut. *Ticks Tick Borne Dis* 10:894–900. <https://doi.org/10.1016/j.ttbdis.2019.04.011>.
 40. Karpathy SE, Slater KS, Goldsmith CS, Nicholson WL, Paddock CD. 2016. *Rickettsia amblyommatis* sp. nov., a spotted fever group Rickettsia associated with multiple species of *Amblyomma* ticks in North, Central and South America. *Int J Syst Evol Microbiol* 66:5236–5243. <https://doi.org/10.1099/ijsem.0.001502>.
 41. Ponnusamy L, Gonzalez A, Van Treuren W, Weiss S, Parobek CM, Juliano

- JJ, Knight R, Roe RM, Apperson CS, Meshnick SR. 2014. Diversity of Rickettsiales in the microbiome of the lone star tick, *Amblyomma americanum*. *Appl Environ Microbiol* 80:354–359. <https://doi.org/10.1128/AEM.02987-13>.
42. McQuiston JH, Zemtsova G, Perniciaro J, Hutson M, Singleton J, Nicholson WL, Levin ML. 2012. Afebrile spotted fever group rickettsia infection after a bite from a Dermacentor variabilis tick infected with Rickettsia montanensis. *Vector Borne Zoonotic Dis* 12:1059–1061. <https://doi.org/10.1089/vbz.2012.1078>.
 43. Carlyon JA, Fikrig E. 2006. Mechanisms of evasion of neutrophil killing by *Anaplasma phagocytophilum*. *Curr Opin Hematol* 13:28–33. <https://doi.org/10.1097/01.moh.0000190109.00532.56>.
 44. Rikihisa Y. 2011. Mechanisms of obligatory intracellular infection with *Anaplasma phagocytophilum*. *Clin Microbiol Rev* 24:469–489. <https://doi.org/10.1128/CMR.00064-10>.
 45. Rikihisa Y. 2015. Molecular pathogenesis of Ehrlichia chaffeensis infection. *Annu Rev Microbiol* 69:283–304. <https://doi.org/10.1146/annurev-micro-091014-104411>.
 46. Ismail N, McBride JW. 2017. Tick-borne emerging infections: ehrlichiosis and anaplasmosis. *Clin Lab Med* 37:317–340. <https://doi.org/10.1016/j.cll.2017.01.006>.
 47. Massung RF, Priestley RA, Miller NJ, Mather TN, Levin ML. 2003. Inability of a variant strain of *Anaplasma phagocytophilum* to infect mice. *J Infect Dis* 188:1757–1763. <https://doi.org/10.1086/379725>.
 48. Massung RF, Courtney JW, Hiratzka SL, Pitzer VE, Smith G, Dryden RL. 2005. *Anaplasma phagocytophilum* in white-tailed deer. *Emerg Infect Dis* 11:1604–1606. <https://doi.org/10.3201/eid1110.041329>.
 49. Tokarz R, Jain K, Bennett A, Briese T, Lipkin WI. 2010. Assessment of polymicrobial infections in ticks in New York state. *Vector Borne Zoonotic Dis* 10:217–221. <https://doi.org/10.1089/vbz.2009.0036>.
 50. Aliota MT, Dupuis AP, Wilczek MP, Peters RJ, Ostfeld RS, Kramer LD, Kramer LD. 2014. The prevalence of zoonotic tick-borne pathogens in Ixodes scapularis collected in the Hudson Valley, New York State. *Vector Borne Zoonotic Dis* 14:245–250. <https://doi.org/10.1089/vbz.2013.1475>.
 51. Adler GH, Levins R. 1994. The island syndrome in rodent populations. *Q Rev Biol* 69:473–490. <https://doi.org/10.1086/418744>.
 52. Stromdahl EY, Nadolny RM, Hickling GJ, Hamer SA, Ogden NH, Casal C, Heck GA, Gibbons JA, Cremeans TF, Pilgard MA. 2018. *Amblyomma americanum* (Acari: Ixodidae) ticks are not vectors of the Lyme disease agent, *Borrelia burgdorferi* (Spirochaetales: Spirochaetales): a review of the evidence. *J Med Entomol* 55:501–514. <https://doi.org/10.1093/jme/tjx250>.
 53. Rosner F, Zarrabi MH, Benach JL, Habicht GS. 1984. Babesiosis in splenectomized adults. Review of 22 reported cases. *Am J Med* 76:696–701. [https://doi.org/10.1016/0002-9343\(84\)90298-5](https://doi.org/10.1016/0002-9343(84)90298-5).
 54. Mah A, Viola GM, Ariza Heredia E, Rezvani K, Kebraiee P, Bhatti MM, Han X, Shpall EJ, Mulanovich VE. 2018. Graft loss attributed to possible transfusion-transmitted ehrlichiosis following cord blood stem cell transplant. *Transpl Infect Dis* 20:e12899. <https://doi.org/10.1111/tid.12899>.
 55. Goel R, Westblade LF, Kessler DA, Sfeir M, Slavinski S, Backenson B, Gebhardt L, Kane K, Laurence J, Scherr D, Bussel J, Dumler JS, Cushing MM, Vasovic LV. 2018. Death from transfusion-transmitted anaplasmosis, New York, USA, 2017. *Emerg Infect Dis* 24:1548–1550. <https://doi.org/10.3201/eid2408.172048>.
 56. Eisen RJ, Eisen L. 2018. The blacklegged tick, *Ixodes scapularis*: an increasing public health concern. *Trends Parasitol* 34:295–309. <https://doi.org/10.1016/j.pt.2017.12.006>.
 57. Dammin GJ, Spielman A, Benach JL, Piesman J. 1981. The rising incidence of clinical Babesia microti infection. *Hum Pathol* 12:398–400. [https://doi.org/10.1016/s0046-8177\(81\)80020-2](https://doi.org/10.1016/s0046-8177(81)80020-2).
 58. Gombert ME, Goldstein EJ, Benach JL, Tenenbaum MJ, Grunwaldt E, Kaplan MH, Eveland LK. 1982. Human babesiosis. Clinical and therapeutic considerations. *JAMA* 248:3005–3007. <https://doi.org/10.1001/jama.1982.03330220049035>.
 59. Grabowski EF, Giardina PJV, Goldberg D, Masur H, Read SE, Hirsch RL, Benach JL. 1982. Babesiosis transmitted by a transfusion of frozen-thawed blood. *Ann Intern Med* 96:466–467. <https://doi.org/10.7326/0003-4819-96-4-446>.
 60. Goethert HK, Telford SR. 2014. Not “out of Nantucket”: Babesia microti in southern New England comprises at least two major populations. *Parasit Vectors* 7:546. <https://doi.org/10.1186/s13071-014-0546-y>.
 61. Lemieux JE, Tran AD, Freemark L, Schaffner SF, Goethert H, Andersen KG, Bazner S, Li A, McGrath G, Sloan L, Vannier E, Milner D, Pritt B, Rosenberg E, Telford S, Bailey JA, Sabeti PC. 2016. A global map of genetic diversity in Babesia microti reveals strong population structure and identifies variants associated with clinical relapse. *Nat Microbiol* 1:16079. <https://doi.org/10.1038/nmicrobiol.2016.79>.
 62. Goethert HK, Molloy P, Berardi V, Weeks K, Telford SR. 2018. Zoonotic Babesia microti in the northeastern U.S.: evidence for the expansion of a specific parasite lineage. *PLoS One* 13:e0193837. <https://doi.org/10.1371/journal.pone.0193837>.
 63. Cavanaugh CE, Muscat PL, Telford SR, Goethert H, Pendlebury W, Elias SP, Robich R, Welch M, Lubelczyk CB, Smith RP. 2017. Fatal deer tick virus infection in Maine. *Clin Infect Dis* 65:1043–1046. <https://doi.org/10.1093/cid/cix435>.
 64. Tavakoli NP, Wang H, Dupuis M, Hull R, Ebel GD, Gilmore EJ, Faust PL. 2009. Fatal case of deer tick virus encephalitis. *N Engl J Med* 360:2099–2107. <https://doi.org/10.1056/NEJMoa0806326>.
 65. Solomon IH, Spera KM, Ryan SL, Helgager J, Andrici J, Zaki SR, Vaitkevicius H, Leon KE, Wilson MR, DeRisi JL, Koo S, Smirnakis SM, De Girolami U. 2018. Fatal Powassan encephalitis (deer tick virus, lineage II) in a patient with fever and orchitis receiving rituximab. *JAMA Neurol* 75:746–750. <https://doi.org/10.1001/jamaneurol.2018.0132>.
 66. Hermance ME, Thangamani S. 2017. Powassan virus: an emerging arbovirus of public health concern in North America. *Vector Borne Zoonotic Dis* 17:453–462. <https://doi.org/10.1089/vbz.2017.2110>.
 67. Frost HM, Schotthoefer AM, Thomm AM, Dupuis AP, Kehl SC, Kramer LD, Fritsche TR, Harrington YA, Knox KK. 2017. Serologic evidence of Powassan virus infection in patients with suspected Lyme disease. *Emerg Infect Dis* 23:1384–1388. <https://doi.org/10.3201/eid2308.161971>.
 68. Krow-Lucal ER, Lindsey NP, Fischer M, Hills SL. 2018. Powassan virus disease in the United States, 2006–2016. *Vector Borne Zoonotic Dis* 18:286–290. <https://doi.org/10.1089/vbz.2017.2239>.
 69. Garcia-Monco JC, Benach JL. 2019. Lyme neuroborreliosis: clinical outcomes, controversy, pathogenesis, and polymicrobial infections. *Ann Neurol* 85:21–31. <https://doi.org/10.1002/ana.25389>.
 70. Mixson TR, Campbell SR, Gill JS, Ginsberg HS, Reichard MV, Schulze TL, Dasch GA. 2006. Prevalence of Ehrlichia, Borrelia, and rickettsial agents in Amblyomma americanum (Acari: Ixodidae) collected from nine states. *J Med Entomol* 43:1261–1268. [https://doi.org/10.1603/0022-2585\(2006\)43\[1261:POEBAR\]2.0.CO;2](https://doi.org/10.1603/0022-2585(2006)43[1261:POEBAR]2.0.CO;2).
 71. Childs JE, Paddock CD. 2003. The ascendancy of Amblyomma americanum as a vector of pathogens affecting humans in the United States. *Annu Rev Entomol* 48:307–337. <https://doi.org/10.1146/annurev.ento.48.091801.112728>.
 72. Sonenshine D. 2018. Range expansion of tick disease vectors in North America: implications for spread of tick-borne disease. *Int J Environ Res Public Health* 15:478. <https://doi.org/10.3390/ijerph15030478>.
 73. Stafford KC, Molaei G, Little EAH, Paddock CD, Karpathy SE, Labonte AM. 2018. Distribution and establishment of the lone star tick in Connecticut and implications for range expansion and public health. *J Med Entomol* 55:1561–1568. <https://doi.org/10.1093/jme/tjy115>.
 74. Jordan RA, Egizi A. 2019. The growing importance of lone star ticks in a Lyme disease endemic county: passive tick surveillance in Monmouth County, NJ, 2006–2016. *PLoS One* 14:e0211778. <https://doi.org/10.1371/journal.pone.0211778>.
 75. Monzón JD, Atkinson EG, Henn BM, Benach JL. 2016. Population and evolutionary genomics of Amblyomma americanum, an expanding arthropod disease vector. *Genome Biol Evol* 8:1351–1360. <https://doi.org/10.1093/gbe/evw080>.
 76. Armstrong PM, Brunet LR, Spielman A, Telford SR. 2001. Risk of Lyme disease: perceptions of residents of a Lone Star tick-infested community. *Bull World Health Organ* 79:916–925.
 77. Beall MJ, Alleman AR, Breitschwerdt EB, Cohn LA, Couto CG, Dryden MW, Guptill LC, Iazbik C, Kania SA, Lathan P, Little SE, Roy A, Saylor KA, Stillman BA, Welles EG, Wolfson W, Yabsley MJ. 2012. Seroprevalence of Ehrlichia canis, Ehrlichia chaffeensis and Ehrlichia ewingii in dogs in North America. *Parasit Vectors* 5:29. <https://doi.org/10.1186/1756-3305-5-29>.
 78. Reeves WK, Loftis AD, Nicholson WL, Czarkowski AG. 2008. The first report of human illness associated with the Panola Mountain Ehrlichia species: a case report. *J Med Case Rep* 2:139. <https://doi.org/10.1186/1752-1947-2-139>.
 79. Regan J, Matthias J, Green-Murphy A, Stanek D, Bertholf M, Pritt BS, Sloan LM, Kelly AJ, Singleton J, McQuiston JH, Hovevar SN, Whittle JP.

2013. A confirmed Ehrlichia ewingii infection likely acquired through platelet transfusion. Clin Infect Dis 56:e105–e107. <https://doi.org/10.1093/cid/cit177>.
80. Esbenshade A, Esbenshade J, Domm J, Williams J, Frangoul H. 2010. Severe ehrlichia infection in pediatric oncology and stem cell transplant patients. Pediatr Blood Cancer 54:776–778. <https://doi.org/10.1002/pcb.22392>.
 81. Nichols Heitman K, Dahlgren FS, Drexler NA, Massung RF, Behravesh CB. 2016. Increasing incidence of ehrlichiosis in the United States: a summary of national surveillance of Ehrlichia chaffeensis and Ehrlichia ewingii infections in the United States, 2008–2012. Am J Trop Med Hyg 94:52–60. <https://doi.org/10.4269/ajtmh.15-0540>.
 82. Varela AS, Luttrell MP, Howerth EW, Moore VA, Davidson WR, Stallknecht DE, Little SE. 2004. First culture isolation of Borrelia lonestari, putative agent of southern tick-associated rash illness. J Clin Microbiol 42:1163–1169. <https://doi.org/10.1128/jcm.42.3.1163-1169.2004>.
 83. Hudman DA, Sargentini NJ. 2016. Detection of Borrelia, Ehrlichia, and Rickettsia spp. in ticks in northeast Missouri. Ticks Tick Borne Dis 7:915–921. <https://doi.org/10.1016/j.ttbdis.2016.04.010>.
 84. Schulze TL, Jordan RA, White JC, Roegner VE, Healy SP. 2011. Geographical distribution and prevalence of selected Borrelia, Ehrlichia, and Rickettsia infections in Amblyomma americanum (Acari: Ixodidae) in New Jersey. J Am Mosq Control Assoc 27:236–244. <https://doi.org/10.2987/11-6111.1>.
 85. James AM, Liveris D, Wormser GP, Schwartz I, Montecalvo MA, Johnson B. 2001. Borrelia lonestari infection after a bite by an Amblyomma americanum tick. J Infect Dis 183:1810–1814. <https://doi.org/10.1086/320721>.
 86. Apperson CS, Engber B, Nicholson WL, Mead DG, Engel J, Yabsley MJ, Dail K, Johnson J, Watson DW. 2008. Tick-borne diseases in North Carolina: is “Rickettsia amblyommii” a possible cause of rickettsiosis reported as Rocky Mountain spotted fever? Vector-Borne Zoonotic Dis 8:597–606. <https://doi.org/10.1089/vbz.2007.0271>.
 87. Weller SJ, Baldrige GD, Munderloh UG, Noda H, Simser J, Kurtti TJ. 1998. Phylogenetic placement of rickettsiae from the ticks Amblyomma americanum and Ixodes scapularis. J Clin Microbiol 36:1305–1317.
 88. Alowaysi M, Chen J, Stark S, Teague K, LaCourse M, Proctor J, Vigil K, Corrigan J, Harding A, Li J, Kurtti T, Zhong J. 2019. Isolation and characterization of a rickettsia from the ovary of a Western black-legged tick, Ixodes pacificus. Ticks Tick Borne Dis 10:918–923. <https://doi.org/10.1016/j.ttbdis.2019.04.017>.
 89. Felsheim RF, Kurtti TJ, Munderloh UG. 2009. Genome sequence of the endosymbiont Rickettsia peacockii and comparison with virulent Rickettsia rickettsii: identification of virulence factors. PLoS One 4:e8361. <https://doi.org/10.1371/journal.pone.0008361>.
 90. Gillespie JJ, Joardar V, Williams KP, Driscoll T, Hostetler JB, Nordberg E, Shukla M, Walenz B, Hill CA, Nene VM, Azad AF, Sobral BW, Caler E. 2012. A Rickettsia genome overrun by mobile genetic elements provides insight into the acquisition of genes characteristic of an obligate intracellular lifestyle. J Bacteriol 194:376–394. <https://doi.org/10.1128/JB.06244-11>.
 91. Benach JL, White DJ, Burgdorfer W, Keelan T, Guirgis S, Altieri RH. 1977. Changing patterns in the incidence of Rocky Mountain spotted fever on Long Island (1971–1976). Am J Epidemiol 106:380–387. <https://doi.org/10.1093/oxfordjournals.aje.a112479>.
 92. Stromdahl EY, Jiang J, Vince M, Richards AL. 2011. Infrequency of Rickettsia rickettsii in Dermacentor variabilis removed from humans, with comments on the role of other human-biting ticks associated with spotted fever group Rickettsiae in the United States. Vector Borne Zoonotic Dis 11:969–977. <https://doi.org/10.1089/vbz.2010.0099>.
 93. Gaines DN, Operario DJ, Stroup S, Stromdahl E, Wright C, Gaff H, Broyhill J, Smith J, Norris DE, Henning T, Lucas A, Houpt E. 2014. Ehrlichia and spotted fever group Rickettsiae surveillance in Amblyomma americanum in Virginia through use of a novel six-plex real-time PCR assay. Vector Borne Zoonotic Dis 14:307–316. <https://doi.org/10.1089/vbz.2013.1509>.
 94. Wormser GP, McKenna D, Scavarda C, Cooper D, El Khoury MY, Nowakowski J, Sudhinda P, Ladenheim A, Wang G, Karmen CL, Demarest V, Dupuis AP, Wong SJ. 2019. Co-infections in persons with early Lyme disease, New York, USA. Emerg Infect Dis 25:748–752. <https://doi.org/10.3201/eid2504.181509>.
 95. Garg K, Meriläinen L, Franz O, Pirttinen H, Quevedo-Diaz M, Croucher S, Gilbert L. 2018. Evaluating polymicrobial immune responses in patients suffering from tick-borne diseases. Sci Rep 8:15932. <https://doi.org/10.1038/s41598-018-34393-9>.
 96. Piesman J, Mather TN, Sinsky RJ, Spielman A. 1987. Duration of tick attachment and Borrelia burgdorferi transmission. J Clin Microbiol 25:557–558.
 97. Katavolos P, Armstrong PM, Dawson JE, Telford SR. 1998. Duration of tick attachment required for transmission of granulocytic ehrlichiosis. J Infect Dis 177:1422–1425. <https://doi.org/10.1086/517829>.
 98. Eisen L. 2018. Pathogen transmission in relation to duration of attachment by Ixodes scapularis ticks. Ticks Tick Borne Dis 9:535–542. <https://doi.org/10.1016/j.ttbdis.2018.01.002>.
 99. Abraham NM, Liu L, Jutras BL, Yadav AK, Narasimhan S, Gopalakrishnan V, Ansari JM, Jefferson KK, Cava F, Jacobs-Wagner C, Fikrig E. 2017. Pathogen-mediated manipulation of arthropod microbiota to promote infection. Proc Natl Acad Sci U S A 114:E781–E790. <https://doi.org/10.1073/pnas.1613422114>.
 100. Levin ML, Schumacher LBM, Snellgrove A. 2018. Effects of Rickettsia amblyommatis infection on the vector competence of Amblyomma americanum ticks for Rickettsia rickettsii. Vector Borne Zoonotic Dis 18:579–587. <https://doi.org/10.1089/vbz.2018.2284>.
 101. Coleman JL, Gebbia JA, Piesman J, Degen JL, Bugge TH, Benach JL. 1997. Plasminogen is required for efficient dissemination of B. burgdorferi in ticks and for enhancement of spirochetemia in mice. Cell 89:1111–1119. [https://doi.org/10.1016/s0092-8674\(00\)80298-6](https://doi.org/10.1016/s0092-8674(00)80298-6).
 102. Liu L, Dai J, Zhao YO, Narasimhan S, Yang Y, Zhang L, Fikrig E. 2012. Ixodes scapularis JAK-STAT pathway regulates tick antimicrobial peptides, thereby controlling the agent of human granulocytic anaplasmosis. J Infect Dis 206:1233–1241. <https://doi.org/10.1093/infdis/jis484>.
 103. Sultana H, Neelakanta G, Kantor FS, Malawista SE, Fish D, Montgomery RR, Fikrig E. 2010. Anaplasma phagocytophilum induces actin phosphorylation to selectively regulate gene transcription in Ixodes scapularis ticks. J Exp Med 207:1727–1743. <https://doi.org/10.1084/jem.20100276>.
 104. Villar M, López V, Ayllón N, Cabezas-Cruz A, López JA, Vázquez J, Alberdi P, de la Fuente J. 2016. The intracellular bacterium Anaplasma phagocytophilum selectively manipulates the levels of vertebrate host proteins in the tick vector Ixodes scapularis. Parasit Vectors 9:467. <https://doi.org/10.1186/s13071-016-1747-3>.
 105. Ceraul SM, Dreher-Lesnick SM, Gillespie JJ, Rahman MS, Azad AF. 2007. New tick defensin isoform and antimicrobial gene expression in response to Rickettsia montanensis challenge. Infect Immun 75:1973–1983. <https://doi.org/10.1128/IAI.01815-06>.
 106. Ceraul SM, Chung A, Sears KT, Popov VL, Beier-Sexton M, Rahman MS, Azad AF. 2011. A Kunitz protease inhibitor from Dermacentor variabilis, a vector for spotted fever group rickettsiae, limits Rickettsia montanensis invasion. Infect Immun 79:321–329. <https://doi.org/10.1128/IAI.00362-10>.
 107. Ginsberg HS. 2008. Potential effects of mixed infections in ticks on transmission dynamics of pathogens: comparative analysis of published records. Exp Appl Acarol 46:29–41. <https://doi.org/10.1007/s10493-008-9175-5>.
 108. Sonenshine DE, Macaluso KR. 2017. Microbial invasion vs. tick immune regulation. Front Cell Infect Microbiol 7:390. <https://doi.org/10.3389/fcimb.2017.00390>.
 109. Shaw DK, Tate AT, Schneider DS, Levashina EA, Kagan JC, Pal U, Fikrig E, Pedra J. 2018. Vector immunity and evolutionary ecology: the harmonious dissonance. Trends Immunol 39:862–873. <https://doi.org/10.1016/j.it.2018.09.003>.
 110. Keirans JE, Durden LA. 1998. Illustrated key to nymphs of the tick genus Amblyomma (Acari: Ixodidae) found in the United States. J Med Entomol 35:489–495. <https://doi.org/10.1093/jmedent/35.4.489>.
 111. Hoskins JD. 1991. Ixodid and argasid ticks. Keys to their identification. Vet Clin North Am Small Anim Pract 21:185–197. [https://doi.org/10.1016/S0195-5616\(91\)50018-8](https://doi.org/10.1016/S0195-5616(91)50018-8).
 112. Dubie TR, Grantham R, Coburn L, Noden BH. 2017. Pictorial key for identification of immature stages of common ixodid ticks found in pastures in Oklahoma. Southwest Entomol 42:1–14. <https://doi.org/10.3958/059.042.0101>.
 113. Tokarz R, Tagliaferro T, Cucura DM, Rochlin I, Sameroff S, Lipkin WI. 2017. Detection of Anaplasma phagocytophilum, Babesia microti, Borrelia burgdorferi, Borrelia miyamotoi, and Powassan Virus in ticks by a multiplex real-time reverse transcription-PCR assay. mSphere 2:e00151-17. <https://doi.org/10.1128/mSphere.00151-17>.
 114. Tokarz R, Kapoor V, Samuel JE, Bouyer DH, Briese T, Lipkin WI. 2009. Detection of tick-borne pathogens by MassTag polymerase chain re-

- action. *Vector Borne Zoonotic Dis* 9:147–152. <https://doi.org/10.1089/vbz.2008.0088>.
115. Schulze TL, Jordan RA, Schulze CJ, Mixson T, Papero M. 2005. Relative encounter frequencies and prevalence of selected *Borrelia*, *Ehrlichia*, and *Anaplasma* infections in *Amblyomma americanum* and *Ixodes scapularis* (Acari: Ixodidae) ticks from central New Jersey. *J Med Entomol* 42:450–456. <https://doi.org/10.1093/jmedent/42.3.450>.
 116. Prusinski MA, Kokas JE, Hukey KT, Kogut SJ, Lee J, Backenson PB. 2014. Prevalence of *Borrelia burgdorferi* (Spirochaetales: Spirochaetaceae), *Anaplasma phagocytophilum* (Rickettsiales: Anaplasmataceae), and *Babesia microti* (Piroplasmida: Babesiidae) in *Ixodes scapularis* (Acari: Ixodidae) collected from recreational lands in the Hudson Valley region, New York State. *J Med Entomol* 51:226–236. <https://doi.org/10.1603/ME13101>.
 117. Steiner FE, Pinger RR, Vann CN, Grindle N, Civitello D, Clay K, Fuqua C. 2007. Infection and co-infection rates of *Anaplasma phagocytophilum* variants, *Babesia* spp., *Borrelia burgdorferi*, and the rickettsial endosymbiont in *Ixodes scapularis* (Acari: Ixodidae) from sites in Indiana, Maine, Pennsylvania, and Wisconsin. *J Med Entomol* 45: 289–297. <https://doi.org/10.1093/jmedent/45.2.289>.
 118. Hamer SA, Roy PL, Hickling GJ, Walker ED, Foster ES, Barber CC, Tsao JI. 2007. Zoonotic pathogens in *Ixodes scapularis*, Michigan. *Emerg Infect Dis* 13:1131–1133. <https://doi.org/10.3201/eid1307.070046>.
 119. Schulze TL, Jordan RA, Healy SP, Roegner VE. 2013. Detection of *Babesia microti* and *Borrelia burgdorferi* in host-seeking *Ixodes scapularis* (Acari: Ixodidae) in Monmouth County, New Jersey. *J Med Entomol* 50:379–383. <https://doi.org/10.1603/me12088>.
 120. Oliver JD, Bennett SW, Beati L, Bartholomay LC. 2017. Range expansion and increasing *Borrelia burgdorferi* infection of the tick *Ixodes scapularis* (Acari: Ixodidae) in Iowa, 1990–2013. *J Med Entomol* 54: 1727–1734. <https://doi.org/10.1093/jme/tjx121>.
 121. Lee X, Coyle DR, Johnson DKH, Murphy MW, McGeehin MA, Murphy RJ, Raffa KF, Paskewitz SM. 2014. Prevalence of *Borrelia burgdorferi* and *Anaplasma phagocytophilum* in *Ixodes scapularis* (Acari: Ixodidae) nymphs collected in managed red pine forests in Wisconsin. *J Med Entomol* 51:694–701. <https://doi.org/10.1603/ME13140>.
 122. Hersh MH, Ostfeld RS, McHenry DJ, Tibbetts M, Brunner JL, Killilea ME, LoGiudice K, Schmidt KA, Keesing F. 2014. Co-infection of blacklegged ticks with *Babesia microti* and *Borrelia burgdorferi* is higher than expected and acquired from small mammal hosts. *PLoS One* 9:e99348. <https://doi.org/10.1371/journal.pone.0099348>.
 123. Feldman KA, Connally NP, Hojgaard A, Jones EH, White JL, Hinckley AF. 2015. Abundance and infection rates of *Ixodes scapularis* nymphs collected from residential properties in Lyme disease-endemic areas of Connecticut, Maryland, and New York. *J Vector Ecol* 40:198–201. <https://doi.org/10.1111/jvec.12153>.
 124. Hutchinson ML, Strohecker MD, Simmons TW, Kyle AD, Helwig MW. 2015. Prevalence rates of *Borrelia burgdorferi* (Spirochaetales: Spirochaetaceae), *Anaplasma phagocytophilum* (Rickettsiales: Anaplasmataceae), and *Babesia microti* (Piroplasmida: Babesiidae) in host-seeking *Ixodes scapularis* (Acari: Ixodidae) from Pennsylvania. *J Med Entomol* 52:693–698. <https://doi.org/10.1093/jme/tjv037>.
 125. Johnson TL, Graham CB, Boegler KA, Cherry CC, Maes SE, Pilgard MA, Hojgaard A, Buttke DE, Eisen RJ. 2017. Prevalence and diversity of tick-borne pathogens in nymphal *Ixodes scapularis* (Acari: Ixodidae) in eastern National Parks. *J Med Entomol* 54:742. <https://doi.org/10.1093/jme/tjw213>.
 126. Johnson TL, Graham CB, Maes SE, Hojgaard A, Fleshman A, Boegler KA, Delory MJ, Slater KS, Karpathy SE, Bjork JK, Neitzel DF, Schiffman EK, Eisen RJ. 2018. Prevalence and distribution of seven human pathogens in host-seeking *Ixodes scapularis* (Acari: Ixodidae) nymphs in Minnesota, USA. *Ticks Tick Borne Dis* 9:1499–1507. <https://doi.org/10.1016/j.ttbdis.2018.07.009>.
 127. Cross ST, Kapuscinski ML, Perino J, Maertens BL, Weger-Lucarelli J, Ebel GD, Stenglein MD. 2018. Co-infection patterns in individual *Ixodes scapularis* ticks reveal associations between viral, eukaryotic and bacterial microorganisms. *Viruses* 10:388. <https://doi.org/10.3390/v10070388>.
 128. Edwards MJ, Russell JC, Davidson EN, Yanusheski TJ, Fleischman BL, Heist RO, Leep-Lazar JG, Stuppi SL, Esposito RA, Suppan LM. 2019. A 4-yr survey of the range of ticks and tick-borne pathogens in the Lehigh Valley region of eastern Pennsylvania. *J Med Entomol* 56: 1122–1134. <https://doi.org/10.1093/jme/tjz043>.