



A *rhII* 5' UTR-Derived sRNA Regulates RhlR-Dependent Quorum Sensing in *Pseudomonas aeruginosa*

Maureen K. Thomason,^a  Maya Voichek,^b Daniel Dar,^{b*} Victoria Addis,^c David Fitzgerald,^c Susan Gottesman,^c Rotem Sorek,^b  E. Peter Greenberg^a

^aDepartment of Microbiology, University of Washington, Seattle, Washington, USA

^bDepartment of Molecular Genetics, Weizmann Institute of Science, Rehovot, Israel

^cLaboratory of Molecular Biology, CCR, NCI, National Institutes of Health, Bethesda, Maryland, USA

ABSTRACT *N*-Acyl homoserine lactone (AHL) quorum sensing (QS) controls expression of over 200 genes in *Pseudomonas aeruginosa*. There are two AHL regulatory systems: the LasR-LasI circuit and the RhlR-RhII system. We mapped transcription termination sites affected by AHL QS in *P. aeruginosa*, and in doing so we identified AHL-regulated small RNAs (sRNAs). Of interest, we noted that one particular sRNA was located within the *rhII* locus. We found that *rhII*, which encodes the enzyme that produces the AHL *N*-butanoyl-homoserine lactone (C4-HSL), is controlled by a 5' untranslated region (UTR)-derived sRNA we name RhIS. We also identified an anti-sense RNA encoded opposite the beginning of the *rhII* open reading frame, which we name asRhIS. RhIS accumulates as wild-type cells enter stationary phase and is required for the production of normal levels of C4-HSL through activation of *rhII* translation. RhIS also directly posttranscriptionally regulates at least one other unlinked gene, *fpvA*. The asRhIS appears to be expressed at maximal levels during logarithmic growth, and we suggest RhIS may act antagonistically to the asRhIS to regulate *rhII* translation. The *rhII*-encoded sRNAs represent a novel aspect of RNA-mediated tuning of *P. aeruginosa* QS.

IMPORTANCE The opportunistic human pathogen *Pseudomonas aeruginosa* possesses multiple quorum sensing systems that regulate and coordinate production of virulence factors and adaptation to different environments. Despite extensive research, the regulatory elements that play a role in this complex network are still not fully understood. By using several RNA sequencing techniques, we were able to identify a small regulatory RNA we named RhIS. RhIS increases translation of RhlII, a key enzyme in the quorum sensing pathway, and represses the *fpvA* mRNA encoding one of the siderophore pyoverdine receptors. Our results highlight a new regulatory layer of *P. aeruginosa* quorum sensing and contribute to the growing understanding of the role regulatory RNAs play in bacterial physiology.

KEYWORDS term-seq, transcriptome, small RNA, Hfq, pyoverdine

The opportunistic pathogen *Pseudomonas aeruginosa*, like many bacteria, has the ability to sense its population density and respond to environmental changes by initiating a gene regulatory system termed quorum sensing (QS). In proteobacteria like *P. aeruginosa*, QS commonly involves diffusible *N*-acylhomoserine lactone (AHL) signaling molecules that are recognized by corresponding transcription factors. When the population density, and thus signal concentration, reaches a critical threshold, a coordinated, population-wide shift in gene expression occurs. This facilitates *P. aeruginosa* adaptation to its environment (1, 2). In *P. aeruginosa*, there are two AHL QS systems: the Las system and the Rhl system. The Las system consists of LasI, which catalyzes synthesis of *N*-3-oxo-dodecanoyl-homoserine lactone (3OC12-HSL), and the

Citation Thomason MK, Voichek M, Dar D, Addis V, Fitzgerald D, Gottesman S, Sorek R, Greenberg EP. 2019. A *rhII* 5' UTR-derived sRNA regulates RhlR-dependent quorum sensing in *Pseudomonas aeruginosa*. *mBio* 10:e02253-19. <https://doi.org/10.1128/mBio.02253-19>.

Editor Carmen Buchrieser, Institut Pasteur

Copyright © 2019 Thomason et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Rotem Sorek, rotem.sorek@weizmann.ac.il, or E. Peter Greenberg, epgreen@uw.edu.

* Present address: Division of Geological and Planetary Science, California Institute of Technology, Pasadena, California, USA.

M.K.T. and M.V. contributed equally to this article.

This article is a direct contribution from Rotem Sorek, a Fellow of the American Academy of Microbiology, who arranged for and secured reviews by Stephen Lory, Harvard Medical School, and Elisabeth Sonnleitner, University of Vienna.

Received 31 August 2019

Accepted 11 September 2019

Published 8 October 2019

3OC12-HSL-dependent transcription factor LasR. The Rhl system consists of RhlI, the *N*-butanoyl-homoserine lactone (C4-HSL) synthase, and the C4-HSL-dependent transcription factor RhlR. Together these two QS circuits activate over 200 genes (3–7). Although much is known about direct control of *P. aeruginosa* genes by LasR and RhlR, less is known about QS-mediated posttranscriptional regulation by noncoding RNA elements.

Over the last two decades, we have learned that noncoding RNA is critical for the posttranscriptional control of gene expression. Posttranscriptional regulation by small regulatory RNAs (sRNAs) is known to occur by two mechanisms: by direct base pairing to target mRNAs or by binding to proteins. In the class of base-pairing sRNAs, *trans*-encoded sRNAs base pair with limited complementarity to target mRNAs mediated by the RNA chaperone protein Hfq and recently discovered ProQ domain-containing proteins (reviewed in reference 8). Antisense RNAs (asRNAs), the second type of base-pairing sRNA, are encoded on the strand opposite to that encoding the mRNA and base pair with the mRNA, generally in the absence of protein chaperones, to regulate either translation or stability of their target mRNA (reviewed in references 9 and 10).

RNA-mediated control of transcription or translation can also involve riboswitches or thermosensors found within the 5' untranslated regions (UTRs) of some genes or operons. They control gene expression or translation by directly binding metabolites or signaling molecules, by sensing pH, or by sensing changes in temperature (11, 12). In the absence of a canonical riboswitch, the secondary structure of the RNA in the 5' UTR can on occasion regulate translation by sequestering access to a ribosome-binding site (RBS). For example, in *P. aeruginosa* the sRNA PhrS regulates expression of *pqsR*, which codes for a transcriptional activator, by binding the RBS of an upstream open reading frame (uORF). The binding alters the RNA structure to activate translation of the uORF, which by translational coupling leads to *pqsR* translation (13). Recent studies suggest riboswitches, 5' UTRs, sRNAs, and asRNAs can regulate each other, thus forming complex regulatory networks (14, 15).

Given the complexity of RNA-based regulation, it is not surprising that sRNAs and QS are intricately linked. For example, similar to other bacteria, *P. aeruginosa* Hfq mutants grow abnormally and are attenuated for virulence (16). In the context of QS, transcript profiling showed that Hfq influences expression of 72 QS-activated genes (16–18). More recently, a high-resolution transcriptome sequencing (RNA-seq) study identified a number of sRNAs, including two that were induced by the LasR QS system (19), while another study identified the sRNA PhrD as a positive regulator of *rhlR* (20). Independent of sRNA-based regulation, two RNA thermometers were shown to control expression of the RhlR-activated *rhlAB* operon, as well as *lasI* (21). Although *P. aeruginosa* genome-wide RNA-seq studies have identified hundreds of potential sRNAs and asRNAs (18, 19, 22–25, 28), none have focused specifically on identifying regulatory RNA elements controlled by QS.

We have used term-seq (26) to quantitatively map 3' ends of RNA in *P. aeruginosa* and identify those ends affected by QS. We did not identify any QS-responsive riboswitches, but we did identify a number of sRNAs not previously associated with QS. There was a strongly QS-induced transcription termination site in the 5' UTR of *rhlI*. Follow-up investigations led us to describe an sRNA we name RhIS, which is derived from the 5' UTR of *rhlI*. RhIS activates translation of *rhlI* and induces C4-HSL production, which can be partially complemented in *trans*. RhIS also acts posttranscriptionally in *trans* to regulate the *fpvA* mRNA, which encodes a siderophore receptor (27). Furthermore, the term-seq analysis revealed an antisense RNA opposite *rhlI*. We call this antisense RNA asRhIS and present evidence that RhIS may act as an asRhIS antagonist in *P. aeruginosa*.

RESULTS

Identification of QS-induced RNA 3' ends by term-seq analysis. We mapped the RNA 3' termini, representing to a large extent the transcription termination sites (TTSs) in a *P. aeruginosa* PAO1 *LasI*, *RhlI* (AHL synthesis) mutant incubated with or without




FIG 1 Term-seq method to identify QS-regulated RNAs. (A) Schematic representation of the experimental setup. (B) Expression and termination of the *lasB* elastase-encoding gene is induced in the presence of AHLs in the PAO1 $\Delta lasI \Delta rhII$ mutant (MPK0493). (C) Expression of the SPA0079 sRNA (28) increases upon addition of AHLs. (D) Expression of the SPA0034 sRNA (28) increases upon addition of AHLs. For panels B to D, the number of reads at the term-seq site represents the average of normalized strand-specific term-seq reads in the dominant term-seq position (see Text S1). Black arrowheads indicate 3' ends with AHLs (+AHLs) measured by term-seq (the directionality of the arrowhead is opposed to the expressed strand), while green ones indicate 3' ends without AHLs (-AHLs). Black lines indicate +AHL RNA-seq reads, and green lines indicate -AHLs RNA-seq reads. Arrows at the bottom of each figure indicate gene annotations, and dashed lines indicate gene annotations that extend beyond that depicted.

added 3OC12-HSL and C4-HSL by using term-seq as diagrammed in Fig. 1A and described in detail in the supplemental material (see Text S1). We identified a total of 804 TTSs associated with annotated *P. aeruginosa* genes or operons (see Table S1, tab A, in the supplemental material).

In addition, we identified 21 RNA termini whose expression levels were elevated or reduced by AHLs (Table 1; Table S1, tab B). We believe this is an underrepresentation of QS-regulated genes because of our stringent analysis criteria and limited experimental conditions. Most of the sites affected by AHLs correspond to the 3' ends of known QS-regulated genes. As an example, term-seq reads depicting the 3' end of *lasB*, which codes for the QS-induced elastase enzyme, were much more abundant in cells incubated with AHLs than in cells without AHLs (Fig. 1B; Table S1, tab B). We also identified AHL-controlled TTSs for a number of previously identified sRNAs not known to be associated with QS (Table 1; Table S1, tab B). Among those regulated by AHLs, two previously identified sRNAs (28) are shown in Fig. 1C and D. Interestingly, one of these sRNAs, designated SPA0034 (28), is located downstream of a known QS-activated gene, PA1869 (5) (Fig. 1D), and might be a 3' UTR-derived sRNA similar to those identified in other bacterial species (29–32). Notably, we identified a premature tran-

TABLE 1 Known small RNAs identified by term-seq as differentially regulated by AHLs

sRNA name	Flanking genes (5'/3')	3' end position ^a	sRNA strand	Fold change ^b	Comments and reference(s)
RhIS/SPA104	<i>rhIR/rhII</i>	3889777	-	55.68	Fig. 2A (28)
SPA0116	PA2768/PA2769	3127925	+	28.56	PA2769, known to be QS regulated (28, 54)
SPA0079	PA2763/PA2764	3123367	-	18.05	Fig. 1C (28)
SPA0034	PA1869/PA1870	2031856	+	3.06	Fig. 1D (28)
SPA0080	PA2789/PA2790	3147657	+	2.94	28
AmiL	<i>amiE</i> /PA3367	3778033	-	2.92	18
pant90	PA0806/PA0807	884182	-	2.26	23

^aIndicates position of the 3' end signal in the PAO1 genome.

^bIndicates fold change with or without AHLs determined as differentially expressed if they changed by more than 2-fold with a *P* value of <0.05.




FIG 2 Analysis of a QS-regulated RNA element in the 5' UTR of *rhII*. (A) RNA-seq and term-seq data for the *rhII* locus in PAO1 $\Delta lasI$ $\Delta rhII$ (MPK0493). Expression and termination of RhIS (orange arrow) located in the 5' UTR of *rhII* increases substantially upon addition of AHLs, producing two alternative isoforms from the same locus. The gap in RNA-seq coverage within *rhII* is due to the deletion of the ORF. The number of reads at the term-seq site (black pillar) was determined as in Fig. 1 (Text S1). The RNA-seq coverage shown is not strand-specific. (B) Northern analysis of RhIS expression in PAO1 $\Delta lasI$ $\Delta rhII$ (MPK0493). Overnight cultures of PAO1 $\Delta lasI$ $\Delta rhII$ were grown as described (Text S1). At an optical density (OD_{600}) of ~ 0.8 , the cultures were split and no AHL or both C4-HSL ($10 \mu M$) and 3OC12-HSL ($2 \mu M$) were added to the cultures. Cells were harvested after 10 and 60 min, RNA was extracted and $10 \mu g$ total RNA was analyzed by Northern blotting with a ^{32}P -labeled oligonucleotide specific to RhIS or 5S as a loading control. RNA from the $\Delta rhIS-I$ (MPK0627) strain collected at OD_{600} of 2.0 was used as a control for band specificity.

scription termination signal in the 5' UTR of *rhII* (Fig. 2). We chose to focus on the AHL-induced *rhII* 5' UTR for two reasons. The expression of this 5' UTR showed a substantial dependence on AHLs (Table 1), and *rhII* codes for the C4-HSL QS signal synthase, which is itself QS activated and required for a full QS response.

Transcription termination in the 5' UTR of *rhII*. The AHL-induced RNA 3'-end in the *rhII* UTR is 34 nucleotides upstream of the *rhII* start codon, and expression of this RNA terminus in AHL-induced cells was over 50 times higher than in uninduced cells (Fig. 2A). Additionally, there is a TTS upstream of *rhII* that maps to the end of the *rhIR* ORF. Thus, the 3' end detected in the 5' UTR of *rhII* is not likely due to transcriptional read-through from *rhIR*. We analyzed whole transcriptome RNA-seq data and found that AHLs induced expression upstream of the *rhII* UTR termination signal by about 100-fold over uninduced cells (Fig. 2A). An overrepresentation of RNA-seq reads in the *rhII* 5' UTR can also be found in previously published RNA-seq data sets of wild-type *P. aeruginosa* strain PA14 (19, 33). We also identified an RNA 3' end in the antisense orientation to the *rhII* ORF overlapping the sequence of *rhII*, although expression of this RNA appeared to be very low (Fig. 3A and see Fig. 6B below).

The 5' UTR does not appear to encode a C4-HSL riboswitch. Because term-seq has been used to discover 5' UTR-derived ribo-regulators that mediate premature transcription termination in other bacteria (26), we asked whether the *rhII* 5' UTR might code for a C4-HSL-responsive riboswitch. We first approached this question by using bioinformatics. Neither the PASIFIC (34) nor RFAM (35) predictive structure analysis programs revealed any putative riboswitch-like motifs in the *rhII* 5' UTR. While informative, these searches are not exhaustive; therefore, we also addressed this question experimentally. We constructed an *E. coli* reporter containing the entire *rhII* 5' UTR through the first 30 codons fused to *lacZ*. Expression of this construct was arabinose inducible. We found that when expression of the construct was activated with arabinose, there was no effect on β -galactosidase levels when C4-HSL was added (see Fig. S1 in the supplemental material). The results from both the bioinformatics and experimental approaches were inconsistent with the idea that the *rhII* 5' UTR is a C4-HSL-responsive riboswitch. We cannot rule out the possibility that this UTR may be responsive to other signaling molecules. However, these results led us to test other possible consequences of early *rhII* transcription termination.




FIG 3 Expression of RhIS as a function of culture growth, LasR, and Hfq. (A) Schematic and sequence of *rhIR-rhIS-rhII* locus. Blue arrow indicates the +1 site of transcription, and red bars indicate the termination points. The *rhIS-rhII* -10 and -35 sequences are underlined and the Las box is boxed in gray. Dashed lines indicate the putative RhIS Rho-independent terminator. (B) Northern analysis of RhIS. WT PAO1 (MPK0409), PAO1 $\Delta lasI \Delta rhII$ (MPK0493), or the RhIS- $\Delta term$ mutant (MPK0555) was grown as described (Text S1). Samples were collected and processed for Northern analysis as in Fig. 2. (C) Influence of LasR and RhIR on RhIS levels. WT PAO1 (MPK0409) and the isogenic PAO1 $\Delta lasR$ (MPK0426) and PAO1 $\Delta rhIR$ (MPK0428) mutants were grown as in panel B, and samples were processed for Northern analysis as in Fig. 2. (D) RhIS levels require Hfq. WT PAO1 (MPK0530) and the isogenic *hfq::aadA* mutant (MPK0529) were grown as in panel B, and samples were processed for Northern analysis as in Fig. 2. In panels B to D, the numbers above the images of autoradiographs indicated the OD₆₀₀ at which cells were harvested for RNA extraction.

The 5' UTR of *rhII* encodes an sRNA. We hypothesized the premature termination of the *rhII* 5' UTR with added AHLs could generate a stable sRNA and investigated this using Northern blot analysis. As shown in Fig. 2B, we detected a QS signal-induced sRNA that is less than 100 nucleotides in length as early as 10 min after exposure to AHLs. This sRNA accumulated for at least 60 min after AHLs were added to the cells. We did not detect the sRNA when we analyzed a 5' UTR-*rhII* deletion mutant, consistent with the conclusion that the sRNA is specific to the *rhII* locus (Fig. 2B). We have named this sRNA RhIS (RhII-associated sRNA). The location and size of RhIS are consistent with a previously identified 70-nucleotide *P. aeruginosa* sRNA (SPA104) that was not known to be AHL induced (28).

We next mapped the 5' end of RhIS by primer extension (see Fig. S2A in the supplemental material) and the 5' end of both RhIS and *rhII* with 5' RACE (rapid amplification of cDNA ends) (Fig. 3A; Fig. S2B). We found the predominant transcript start site of both RhIS and *rhII* corresponded to the +1 position of *rhII* transcription previously reported (36). We also identified a stem-loop structure followed by a polyuridine tract, consistent with a Rho-independent transcriptional terminator sequence immediately upstream of the RNA 3' end detected in the *rhII* 5' UTR sequence (Fig. 3A). By using the boundaries defined by the transcript start site and term-seq, we infer that RhIS is about 70 nucleotides in length. A transcript of this size is consistent

with the RhIS band seen by Northern analysis (Fig. 2B). Our data suggest that, in the presence of AHLs, the *rhII* locus can be transcribed from a single *rhII* promoter into two isoforms: the long isoform encoding full-length *rhII* mRNA and, as our RNA-seq data suggest, a more abundant short RhIS resulting from premature transcription termination within the 5' UTR (Fig. 2A).

RhIS is regulated by QS and is dependent on Hfq. We next examined RhIS production in wild-type *P. aeruginosa*. Northern blotting showed that RhIS levels in early-logarithmic-phase cells were relatively low, increased in late logarithmic phase, and were at maximal levels in stationary-phase cells (Fig. 3B). There is a LasR binding site in the promoter region of *rhII*, and *rhII* is activated strongly by LasR and weakly by RhIR (36). Because the transcript starts for RhIS and *rhII* appeared to be the same, we hypothesized that RhIS transcription would be activated primarily by LasR and to a lesser extent by RhIR. To test this hypothesis, we monitored RhIS levels in strains deleted for *lasR* and *rhIR* by Northern blotting. As predicted, RhIS levels in the LasR mutant were very low and RhIS was modestly decreased in the RhIR mutant (Fig. 3C). Thus, the increase in RhIS as a function of growth appears to be primarily a consequence of LasR-dependent QS induction.

A previous report showed that there was a marginal effect of Hfq on *rhII* mRNA levels (17). We asked whether Hfq might affect RhIS and the *rhII* transcript differently. In fact, Northern blotting showed very low levels of RhIS in an Hfq mutant in comparison to levels in wild-type cells (Fig. 3D). This is consistent with the conclusion that the two transcripts produced from the *rhl* locus have different requirements for Hfq. RhIS levels are drastically altered in the absence of Hfq, while *rhII* levels are only slightly altered (~1.5-fold decreased in an *hfq* mutant) (17).

Disruption of the RhIS terminator reduces C4-HSL production. From the term-seq analysis, we estimate that RhIS is the predominant transcript derived from the *rhII* promoter, with full-length *rhII* mRNA accounting for between 3 and 15% of the total at steady state. RNA structure prediction using Mfold (37) showed a predicted structure with a 5' end hairpin and a Rho-independent terminator (Fig. 4A). We hypothesized that disruption of the Rho-independent terminator should result in increased transcriptional readthrough producing more full-length *rhII* transcript and thus elevated levels of C4-HSL. We tested this hypothesis by constructing a strain where the Rho-independent terminator is deleted (RhIS- Δ term). Surprisingly, the amount of *rhII* mRNA in this mutant was similar to that in wild-type cells (see Fig. S3 in the supplemental material), but RhIS levels were lower (Fig. 3B). The deletion may have eliminated an essential Hfq binding site present in the terminator (38). The terminator deletion mutant also produced almost 10-fold less C4-HSL than the wild-type (Fig. 4B). Because the wild type and mutant had roughly equivalent levels of *rhII* mRNA, we surmise that the defect in C4-HSL production in the terminator mutant is due to RhIS-mediated posttranscriptional regulation of *rhII*.

To gain insight into whether the terminator structure or sequence of RhIS was responsible for the phenotype exhibited by the terminator deletion mutant, we made point mutations in the RhIS sequence. These point mutations (Fig. 4A, Mut A) should disrupt the terminator structure while leaving the sequence largely intact. We also made a compensatory mutation that should restore base pairing in the terminator and thus recover the structure (Fig. 4A, Mut A+B). The structure-disrupting mutant and the compensating mutant had similar levels of *rhII* mRNA but produced low levels of C4-HSL and low levels of RhIS (Fig. 4B to D). These data are consistent with the conclusion that the sequence of the RNA in the RhIS terminator region is essential for RhIS function, and RhIS is important for C4-HSL production but not *rhII* mRNA levels. It is conceivable that Hfq binds to RhIS in this region to promote the sRNA-mediated posttranscriptional regulation of *rhII*.

The defect in C4-HSL production can be partially restored by *trans*-complementation of RhIS on a multicopy plasmid (pRhIS) in the RhIS terminator structure mutation strain (Fig. 4E). Arabinose-induced *P. aeruginosa* (pRhIS) had RhIS transcript levels similar to




FIG 4 RhIS regulation of RhII translation and C4-HSL production. (A) Predicted Mfold (37) structure of the RhIS sRNA. Gray boxes indicate positions and red letters indicate nucleotides changed for MutA, MutA+B and MutC point mutants. Cyan brackets indicate nucleotides deleted in the RhIS- Δterm mutant. (B) C4-HSL levels in RhIS point mutant strains compared to wild type. Supernatant was collected from overnight cultures of WT PAO1 (MPK0409) and the isogenic RhIS- Δterm mutant (MPK0555), MutA mutant (MPK0576), MutA+B mutant (MPK0619), and as a control, the PAO1 $\Delta\text{lasI}\Delta\text{rhII}$ mutant (MPK0493) after 24 h. C4-HSL was extracted and measured by using the C4-HSL bioassay. Values are the means of three biological and two technical replicates, and error bars are standard deviations. (C) Levels of *rhII* mRNA in the wild-type PAO1, MutA, and MutA+B strains were determined by quantitative reverse transcription-PCR (qRT-PCR) using primers specific to the *rhII* open reading frame. The amount of *rhII* mRNA was calculated with a standard curve and normalized to levels of the housekeeping control gene *groEL*. Values are the mean of two biological and two technical replicates. Bars are standard deviations. (D) Levels of RhIS in the point mutant strains. The wild type and the isogenic MutA and MutA+B mutants were grown and Northern analysis performed as in Fig. 3. (E) RhIS can partially complement the MutA C4-HSL production defect when provided in *trans*. Wild-type PAO1 and the PAO1 MutA point mutant were transformed with the empty pMKT1 vector or pRhIS. Single colonies were inoculated into LB plus 50 mM MOPS and grown in the presence or absence of 1% L-arabinose at 37°C. After 24 h, C4-HSL was extracted and measured. Results are means of three biological and two technical replicates, and bars are the standard deviation. (F) Western blot analysis of VSV-G epitope-tagged RhII. Strains: WT-RhII-VSV-G (MPK0698), MutA-VSV-G (MPK0689), MutA+B-VSV-G (MPK0697), asRhIS-1-VSV-G (MPK0687), and $\Delta\text{rhIS}\Delta\text{rhII}$ (MPK0627). Cells were grown as in panel B. RhII levels were normalized to the corresponding RNA polymerase band and are presented as a percentage of wild type. The image is representative of three independent experiments.

those of wild-type cells indicating expression from the pRhIS plasmid is physiologically relevant (see Fig. S4A in the supplemental material). Additionally, when an arabinose-induced promoter-*rhII* fusion was integrated in the *Escherichia coli* chromosome, the cells produced micromolar amounts of C4-HSL similar to those of wild-type *P. aeruginosa* (Fig. S4B). This indicates the mechanism by which RhIS induces *rhII* translation is either conserved between *E. coli* and *P. aeruginosa* or contained entirely within the RhIS-*rhII* locus. Our data suggest that high levels of RhIS are required to maintain wild-type levels of C4-HSL but are not required for *rhII* mRNA accumulation. In our experiments, *rhII* mRNA levels remain unchanged regardless of RhIS expression, while C4-HSL levels vary in mutants expressing less RhIS.

RhIS controls translation of *rhII* mRNA. Because mutants that produce normal levels of *rhII* mRNA but low levels of RhIS produce low levels of C4-HSL, we hypothesized that RhIS might stimulate translation of *rhII* mRNA. To test this hypothesis, we incorporated a C-terminal vesicular stomatitis virus glycoprotein G (VSV-G) epitope tag at the native *rhII* locus such that it encoded an RhII-VSV-G polypeptide. We examined levels of this tagged RhII in our RhIS mutant strains by Western blotting (Fig. 4F). Consistent with the hypothesis, RhII levels were lower in the RhIS mutants than in the wild type. We confirmed that RhII-VSV-G was active by showing the RhII-VSV-G-tagged version produced C4-HSL levels comparable to those of the native RhII protein (see Fig. S5 in the supplemental material). Thus, RhIS appears to affect translation but not the *rhII* mRNA.

Direct regulation of *fpvA* by RhIS. It seemed possible that RhIS might affect translation of genes other than *rhII*. To assess this possibility, we used the base-pairing prediction algorithm TargetRNA2 (39) to search for genomic regions that might base pair with RhIS (Table S1, tab C). The hit with the most extensive base-pairing complementarity was a 16 nucleotide region in the first five codons of the *fpvA* open reading frame that pairs with RhIS (Fig. 5A). The *fpvA* gene product is a receptor for the *P. aeruginosa* siderophore pyoverdine (27).

To test the hypothesis that RhIS controls *fpvA* expression, we created an arabinose-inducible translational reporter containing the *fpvA* 5' UTR through the first 25 codons of the *fpvA* ORF fused in frame to *lacZ* and placed this construct on the chromosome of *E. coli*. In this construct, we either expressed RhIS on an arabinose-inducible plasmid (pRhIS) or included the empty vector (pMKT1). Levels of β -galactosidase in arabinose-grown cells containing pRhIS were about 25% of the levels in cells without RhIS (Fig. 5B). To test whether the RhIS repression of *fpvA* was by direct base pairing via the 16-nucleotide region identified in the TargetRNA2 analysis, we did the following: We first changed the RhIS sequence to disrupt base pairing with *fpvA*. When we used a plasmid (pMutC) expressing the mutant RhIS in place of wild-type RhIS, *fpvA-lacZ* expression was not repressed (Fig. 5B). We then constructed an arabinose-inducible *fpvA-1-lacZ* reporter with a mutation that compensated for the pMutC mutation, and this restored *lacZ* repression (Fig. 5C). We note that levels of *fpvA-1-lacZ* expression in cells containing the compensatory mutation are lower than those in cells containing the wild-type *fpvA-lacZ* fusion. We believe this may be due to the fact that changing the sequence of the first few *fpvA* codons decreased translation efficiency. These experiments indicate that RhIS can serve to regulate *fpvA* mRNA translation by a direct base-pairing mechanism, and they point to a link between QS and iron homeostasis in *P. aeruginosa*.

The *rhII* antisense RNA may be involved in regulating *rhII* translation. As mentioned earlier, we detected low levels of an *rhII* antisense RNA in our term-seq analysis (Fig. 3A). A map showing the chromosomal region encoding this asRNA with the term-seq detected TTS is shown in Fig. 6A. We confirmed the existence of the antisense RNA, which we call asRhIS, by Northern blot analysis. Expression of asRhIS was at peak, although still low, abundance during logarithmic growth of *P. aeruginosa* (Fig. 6B). The size of the asRhIS band is slightly less than 200 nucleotides. Because asRhIS is in low abundance, we were unable to map the exact transcription start site by




FIG 5 RhIS regulation of FpvA production. (A) TargetRNA2 (39) RhIS and *fpvA* base-pairing predictions. The lines over and under the sequences show mutations we constructed and are relative to the start codon for *fpvA* and the +1 site of RhIS transcription. (B) Negative regulation of *fpvA-lacZ* by RhIS. The reporter strain PM1205 *fpvA-lacZ* (MPK0704) was transformed with the control vector pMKT1, pRhIS, or the pMutC mutant derivative. β -Galactosidase levels were assayed after 3 h of induction with 0.4% arabinose. The averages of three independent assays are shown, and error bars are standard deviations. (C) Disruption and restoration of base pairing between RhIS and *fpvA*. The pMKT1 vector, pRhIS or pMutC plasmids were transformed into the PM1205 *fpvA-1-lacZ* mutant strain (MPK0712), which carries compensatory mutations to restore regulation to MutC. β -Galactosidase levels were assayed as in panel B.

5' RACE. However, based on the TTS determined by our term-seq analysis and the size of the band on the Northern blot, we identified a putative TSS (transcription start site) for asRhIS (Fig. 6A; see Fig. S6 in the supplemental material). We predict asRhIS overlaps the beginning of the *rhII* ORF as well as the entire *rhII* 5' UTR. Previous transcription start site mapping identified an antisense TSS in strain *P. aeruginosa* PA14 in close proximity (Fig. 6A), ~40 nucleotides upstream of our putative TSS (19).

We could not identify a σ^{70} promoter-like -10 and -35 region for either the putative PAO1 asRhIS +1 or the known PA14 +1, but we did identify a possible promoter consistent with a *P. aeruginosa* extracytoplasmic function (ECF) σ^E consensus (Fig. 6A) upstream of the putative TSS of the asRhIS in PAO1 (40). Surprisingly, disrupting this sequence increased asRhIS expression (Fig. 6B), leading to decreased C4-HSL (Fig. 6C) and about 25% less RhII protein compared to the wild type (Fig. 4F). *P. aeruginosa* has 19 ECF sigma factors (reviewed in reference 41). We do not know which of these might be involved in asRhIS induction, and we leave it to future studies to elucidate whether and how asRhIS regulates *rhII*.

DISCUSSION

By term-seq mapping of *P. aeruginosa* TTSs in a QS AHL signal synthesis mutant with or without added AHLs, we identified a number of QS-regulated sRNAs. Our list includes seven AHL-regulated sRNAs, all of which had been detected previously but never reported to be associated with QS (Table 1). Three of these sRNAs were highly induced by AHLs (Table 1, Fig. 1C and D, and Fig. 2A) while the other four showed weaker




FIG 6 An antisense RNA is encoded in the *rhII* locus and regulates C4-HSL production. (A) Schematic and sequence of the *asRhIS* promoter region, including the overlap with *RhIS* and *rhII*. Blue boxes indicate the putative +1 of transcription for PAO1 and the known +1 for PA14 and red box indicates the *asRhIS* termination point. The dashed line indicates a predicted σ^E site for *asRhIS* and the consensus sequence is indicated above. Red letters are the nucleotide changes for the *asRhIS-1* mutant. For reference and orientation the *rhII* start codon, *rhIS* start and stop, and *rhIR* stop codon are indicated. The ribosome binding site (RBS) for *rhII* is shaded gray. (B) The *asRhIS-1* mutation increases expression of *asRhIS*. WT PAO1 and the isogenic *asRhIS-1* promoter mutant (MPK0637) were grown and processed for Northern analysis as in Fig. 3 with an oligonucleotide specific to *asRhIS*. * indicates the wild-type *asRhIS* transcript band. (C) C4-HSL levels are reduced in the *asRhIS-1* promoter strain. Wild-type PAO1 (MPK0409) and *asRhIS-1* (MPK0637) were grown and C4-HSL levels were determined as in Fig. 4. Values are means of three biological and two technical replicates for each strain, and error bars are standard deviations. ** indicates $P < 0.005$ using an unpaired *t* test with Welch's correction.

induction by AHLs. In our previous high-resolution RNA-seq analysis of *P. aeruginosa*, we described two LasR-activated sRNAs, Lrs1 and Lrs2 (19), which were not identified in our term-seq analysis. We stress that our analysis was not an exhaustive mapping of QS-dependent sRNA expression. We analyzed only one *P. aeruginosa* strain grown under one condition, 60 min following exposure to AHLs. Furthermore, we used stringent requirements to call a TTS. However, our analysis opens an avenue for future discovery of *P. aeruginosa* QS-dependent sRNA expression.

Here we focused on the most highly AHL-induced sRNA under our conditions, which we have called *RhIS*. *RhIS* is encoded in the 5' UTR of the C4-HSL synthesis *rhII* gene, it is 70 nucleotides in length, it appears to require Hfq, it stimulates *rhII* mRNA translation in a *trans*-acting fashion, and it interferes with the posttranscriptional regulation of an unlinked gene, which codes for the pyoverdine receptor FpvA. Although *RhIS* was induced when we added both 3OC12-HSL and C4-HSL to growing cells, we presume that induction is primarily a response to 3OC12-HSL because *RhIS* expression showed a strong dependence on the 3OC12-HSL receptor LasR and only a weak dependence on the C4-HSL receptor RhIR.

The RhIS and *rhII* transcript start sites appear to be one and the same: There are two RNA isoforms produced from this transcription start site, the shorter RhIS and the longer *rhIS-rhII* isoforms. Some sRNAs can repress premature transcription termination within a 5' UTR by binding and inhibiting Rho-dependent termination (42). However, the presence of a Rho-independent terminator at the 3' end of RhIS (Fig. 3A and 4A) suggests that the long *rhIS-rhII* isoform might result from leaky or imperfect Rho-independent transcription termination rather than by inhibition of Rho-dependent termination. Whether the efficiency of *rhIS-rhII* transcription termination changes upon different growth or stress conditions, as described previously for the *E. coli* SgrS and RybB sRNAs (43), is unclear at this point. However, similar instances of transcriptional readthrough have been shown in other bacteria. For example, in *Salmonella*, leaky transcriptional readthrough of the *IrsK* sRNA terminator leads to a long *IrsK-orf45-anrP* transcript and a stable short *IrsK* sRNA, which can then act in *trans* to increase translation of *orf45* and *anrP* (44). Similarly, incomplete transcription termination of the *Salmonella gltIJKL* operon, which codes for a glutamate-aspartate transporter, produces either the long *gltIJKL* mRNA or a short *gltI* mRNA from which the Hfq-dependent SroC sRNA is processed (45). The SroC sRNA then acts as a sponge to relieve repression of the *gltIJKL* operon. For reasons discussed below, we hypothesize that RhIS may be functioning to relieve interference with *rhII* mRNA translation perhaps by titration of the asRhIS by a SroC sponge-like mechanism.

We find it interesting that the RhIS isoform but not the full-length isoform required Hfq for their function. It is not surprising that the long isoform does not require Hfq, as *rhII* transcript levels are minimally altered in an Hfq mutant (17). This sort of differential response of an sRNA and a longer RNA containing the sRNA sequence to Hfq is not unique to RhIS and *rhII*. Recently it was shown in *E. coli* the Rho-independent terminators in the Hfq dependent sRNAs SgrS and RyhB allowed transcriptional readthrough, which produces longer Hfq-independent transcripts (43).

Previous work showed that despite minimal changes in the *rhII* mRNA in the absence of Hfq, translation of *rhII* and C4-HSL production were reduced in an Hfq mutant (17). We can now explain these reductions. We show that Hfq is important for RhIS, and RhIS is required for normal *rhII* translation and thus C4-HSL production. We note that regulation of *rhII* is complex and also affected by sRNAs other than RhIS. The RNA binding protein RsmA, which binds GGA motifs in the loops of RNA hairpins to repress translation, was shown to repress *rhII* translation and C4-HSL production (46). Additionally, the sRNA RsmY, which binds to RsmA to relieve translational repression, has also been implicated in *rhII* regulation and C4-HSL production by an indirect mechanism involving Hfq stabilization of RsmY (17). Although several GGA motifs are present in RhIS, most are not present in the predicted hairpin loops (although one GGA motif is partially buried), suggesting RhIS likely does not affect translation of *rhII* through titration of RsmA.

We show that RhIS can function to stimulate *rhII* translation in a *trans*-acting fashion. RhIS expressed from a multicopy plasmid (at physiologically relevant levels) restores C4-HSL levels almost to those of wild-type PAO1 (Fig. 4E; Fig. S4A). This finding rules out the possibility the regulation is due to inherent factors in the *rhIS-rhII* transcript itself (e.g., RNA structure alters transcription or translation efficiency of *rhII*). What is unclear at this point is how RhIS mediates this positive regulation of *rhII*. We have two models for how this RhIS regulation may occur. First RhIS could act directly on *rhII* to relieve translational repression mediated by a highly structured 5' UTR. Mfold predicts the secondary structure of the *rhII* 5' UTR occludes the primary RBS by a stem-loop structure (see Fig. S7 in the supplemental material). It is possible RhIS activates *rhII* in *trans* by base pairing to sequences opposite the *rhII* RBS, relieving occlusion and facilitating translation. Examples of this type of regulation exist in *Pseudomonas* (and other bacteria) where an sRNA base pairs to a highly structured 5' UTR to relieve translation repression mediated by RBS occlusion (13, 47–49). Alternatively, as we propose below, RhIS could act in *trans* by a sponge-like mechanism to sequester the asRhIS that appears to repress *rhII* translation.

Our term-seq and RNA-seq analyses also uncovered the antisense RNA asRhIS. The existence of this RNA was confirmed by Northern blotting, and the Northern blotting also revealed that while expression of the asRhIS is low compared to that of RhIS, it is detectable in early-logarithmic-phase cells and not in stationary-phase cells (Fig. 6B). The asRhIS overlaps the beginning of the *rhII* ORF, RBS, and the 5' UTR of *rhII*. It is possible that asRhIS base pairs with the *rhII* mRNA to block translation. Given these data, it is possible that RhIS acts as an asRhIS sponge due to the extensive predicted complementarity between RhIS and asRhIS. Under this hypothesis, this potential interaction between RhIS and asRhIS would sequester asRhIS and relieve the translational repression of *rhII*. Although this is an intriguing hypothesis, the relationship between RhIS and asRhIS requires further investigation.

Finally, besides having a role in *rhII* autoregulation, we identified a region of RhIS complementarity in the mRNA of the pyoverdine receptor gene *fpvA*. By analyzing the influence of mutations in RhIS and compensatory mutations in the *fpvA* 5' UTR on a *fpvA-lacZ* translational fusion we showed that RhIS interferes with translation of *fpvA* via a direct base-pairing mechanism. We have not investigated the physiological significance of this interaction, nor have we searched exhaustively for other potential RhIS-regulated mRNAs; however, we have provided our TargetRNA2 list as a resource for the community (Table S1, tab C). We have thus described an additional layer of gene regulation in the intricate *P. aeruginosa* quorum sensing circuitry.

MATERIALS AND METHODS

Details of additional materials and methods are provided in Text S1.

Bacteria and growth conditions. The bacterial strains, plasmids, and oligonucleotides used in this study are described in Table S1, tab D. Details of strain construction and experimental growth conditions are listed in the Text S1.

RNA extraction, library preparation and sequencing. For RNA-seq, the RNA was extracted using TRIzol and phenol-chloroform. Whole transcriptome RNA-seq libraries and term-seq libraries were prepared as described previously (26).

RNA extraction for Northern blotting. RNA extraction for northern analysis was performed by hot acid phenol-chloroform extraction as described previously with minor changes (see Text S1) (50).

Northern blot analysis. Northern blotting was performed as described previously (50) with minor modifications. Briefly, RNA was separated on 8% polyacrylamide–6 M urea gel (National Diagnostics) and transferred to a Hybond-XL membrane (GE Healthcare). Membranes were probed with [³²P]ATP end-labeled oligonucleotides specific to the desired transcript (Table S1, tab D) and exposed to Amersham Hyperfilm MP (GE-Healthcare) at –80°C.

C4-HSL and β-galactosidase measurements. C4-HSL was ethyl acetate extracted from 24-h LB plus 50 mM MOPS (morpholinepropanesulfonic acid) culture supernatant as described previously (51). The amount of C4-HSL was determined by using an *E. coli* (pECP61.5) bioassay (52, 53) and the Tropix Galacto-Light Plus reagent (Invitrogen).

Data availability. RNA-seq and term-seq data sets have been deposited in the European Nucleotide Database (ENA) under study accession no. [PRJEB31965](https://doi.org/10.1101/2019.08.28.274848).

SUPPLEMENTAL MATERIAL

Supplemental material for this article may be found at <https://doi.org/10.1128/mBio.02253-19>.

TEXT S1, PDF file, 0.3 MB.

FIG S1, PDF file, 0.2 MB.

FIG S2, PDF file, 0.2 MB.

FIG S3, PDF file, 0.3 MB.

FIG S4, PDF file, 0.1 MB.

FIG S5, PDF file, 0.1 MB.

FIG S6, PDF file, 0.2 MB.

FIG S7, PDF file, 0.2 MB.

TABLE S1, XLSX file, 0.6 MB.

ACKNOWLEDGMENTS

We thank M. Schuster, A. Dandakar, P. Mandin, E. Fozo, and H. Schweizer for providing strains and plasmids.

This work was supported by U.S. Public Health Service (USPHS) grant GM59026 to E.P.G. and European Research Council (ERC) grant ERC-CoG 681203 to R.S. Support for

M.K.T. was provided by a Cystic Fibrosis Foundation Postdoctoral Fellowship. M.V. is a Clore Scholar and was supported by the Clore Israel Foundation.

REFERENCES

- Schuster M, Sexton DJ, Diggle SP, Greenberg EP. 2013. Acyl-homoserine lactone quorum sensing: from evolution to application. *Annu Rev Microbiol* 67:43–63. <https://doi.org/10.1146/annurev-micro-092412-155635>.
- Winstanley C, O'Brien S, Brockhurst MA. 2016. *Pseudomonas aeruginosa* evolutionary adaptation and diversification in cystic fibrosis chronic lung infections. *Trends Microbiol* 24:327–337. <https://doi.org/10.1016/j.tim.2016.01.008>.
- Chugani S, Kim BS, Phattarasukol S, Brittnacher MJ, Choi SH, Harwood CS, Greenberg EP. 2012. Strain-dependent diversity in the *Pseudomonas aeruginosa* quorum-sensing regulon. *Proc Natl Acad Sci U S A* 109: E2823–E2831. <https://doi.org/10.1073/pnas.1214128109>.
- Schuster M, Greenberg EP. 2007. Early activation of quorum sensing in *Pseudomonas aeruginosa* reveals the architecture of a complex regulon. *BMC Genomics* 8:287. <https://doi.org/10.1186/1471-2164-8-287>.
- Schuster M, Lostroh CP, Ogi T, Greenberg EP. 2003. Identification, timing, and signal specificity of *Pseudomonas aeruginosa* quorum-controlled genes: a transcriptome analysis. *J Bacteriol* 185:2066–2079. <https://doi.org/10.1128/jb.185.7.2066-2079.2003>.
- Wagner VE, Bushnell D, Passador L, Brooks AI, Iglewski BH. 2003. Microarray analysis of *Pseudomonas aeruginosa* quorum-sensing regulons: effects of growth phase and environment. *J Bacteriol* 185:2080–2095. <https://doi.org/10.1128/jb.185.7.2080-2095.2003>.
- Whiteley M, Lee KM, Greenberg EP. 1999. Identification of genes controlled by quorum sensing in *Pseudomonas aeruginosa*. *Proc Natl Acad Sci U S A* 96:13904–13909. <https://doi.org/10.1073/pnas.96.24.13904>.
- Olejniczak M, Storz G. 2017. ProQ/FinO-domain proteins: another ubiquitous family of RNA matchmakers? *Mol Microbiol* 104:905–915. <https://doi.org/10.1111/mmi.13679>.
- Georg J, Hess WR. 2011. *cis*-antisense RNA, another level of gene regulation in bacteria. *Microbiol Mol Biol Rev* 75:286–300. <https://doi.org/10.1128/MMBR.00032-10>.
- Thomason MK, Storz G. 2010. Bacterial antisense RNAs: how many are there, and what are they doing? *Annu Rev Genet* 44:167–188. <https://doi.org/10.1146/annurev-genet-102209-163523>.
- Sherwood AV, Henkin TM. 2016. Riboswitch-mediated gene regulation: novel RNA architectures dictate gene expression responses. *Annu Rev Microbiol* 70:361–374. <https://doi.org/10.1146/annurev-micro-091014-104306>.
- Kortmann J, Narberhaus F. 2012. Bacterial RNA thermometers: molecular zippers and switches. *Nat Rev Microbiol* 10:255–265. <https://doi.org/10.1038/nrmicro2730>.
- Sonnleitner E, Gonzalez N, Sorger-Domenigg T, Heeb S, Richter AS, Backofen R, Williams P, Hüttenhofer A, Haas D, Bläsi U. 2011. The small RNA PhrS stimulates synthesis of the *Pseudomonas aeruginosa* quorum-sensing signal. *Mol Microbiol* 80:868–885. <https://doi.org/10.1111/j.1365-2958.2011.07620.x>.
- Johansson J, Mandin P, Renzoni A, Chiaruttini C, Springer M, Cossart P. 2002. An RNA thermosensor controls expression of virulence genes in *Listeria monocytogenes*. *Cell* 110:551–561. [https://doi.org/10.1016/s0092-8674\(02\)00905-4](https://doi.org/10.1016/s0092-8674(02)00905-4).
- Loh E, Dussurget O, Gripenland J, Vaitkevicius K, Tiensuu T, Mandin P, Repoila F, Buchrieser C, Cossart P, Johansson J. 2009. A *trans*-acting riboswitch controls expression of the virulence regulator PrfA in *Listeria monocytogenes*. *Cell* 139:770–779. <https://doi.org/10.1016/j.cell.2009.08.046>.
- Sonnleitner E, Hagens S, Rosenau F, Wilhelm S, Habel A, Jäger KE, Blasi U. 2003. Reduced virulence of a *hfr* mutant of *Pseudomonas aeruginosa* O1. *Microb Pathog* 35:217–228. [https://doi.org/10.1016/S0882-4010\(03\)00149-9](https://doi.org/10.1016/S0882-4010(03)00149-9).
- Sonnleitner E, Schuster M, Sorger-Domenigg T, Greenberg EP, Bläsi U. 2006. Hfq-dependent alterations of the transcriptome profile and effects on quorum sensing in *Pseudomonas aeruginosa*. *Mol Microbiol* 59: 1542–1558. <https://doi.org/10.1111/j.1365-2958.2006.05032.x>.
- Sonnleitner E, Sorger-Domenigg T, Madej MJ, Findeiss S, Hackermüller J, Hüttenhofer A, Stadler PF, Bläsi U, Moll I. 2008. Detection of small RNAs in *Pseudomonas aeruginosa* by RNomics and structure-based bioinformatic tools. *Microbiology* 154:3175–3187. <https://doi.org/10.1099/mic.0.2008/019703-0>.
- Wurtzel O, Yoder-Himes DR, Han K, Dandekar AA, Edelhelt S, Greenberg EP, Sorek R, Lory S. 2012. The single-nucleotide resolution transcriptome of *Pseudomonas aeruginosa* grown in body temperature. *PLoS Pathog* 8:e1002945. <https://doi.org/10.1371/journal.ppat.1002945>.
- Malgaonkar A, Nair M. 2019. Quorum sensing in *Pseudomonas aeruginosa* mediated by RhlR is regulated by a small RNA PhrD. *Sci Rep* 9:432. <https://doi.org/10.1038/s41598-018-36488-9>.
- Grosso-Becerra MV, Croda-García G, Merino E, Servín-González L, Mojica-Espinosa R, Soberón-Chávez G. 2014. Regulation of *Pseudomonas aeruginosa* virulence factors by two novel RNA thermometers. *Proc Natl Acad Sci U S A* 111:15562–15567. <https://doi.org/10.1073/pnas.1402536111>.
- Ferrara S, Carloni S, Fulco R, Falcone M, Macchi R, Bertoni G. 2015. Post-transcriptional regulation of the virulence-associated enzyme AlgC by the sigma(22)-dependent small RNA ErsA of *Pseudomonas aeruginosa*. *Environ Microbiol* 17:199–214. <https://doi.org/10.1111/1462-2920.12590>.
- Gomez-Lozano M, Marvig RL, Molin S, Long KS. 2012. Genome-wide identification of novel small RNAs in *Pseudomonas aeruginosa*. *Environ Microbiol* 14:2006–2016. <https://doi.org/10.1111/j.1462-2920.2012.02759.x>.
- Gomez-Lozano M, Marvig RL, Molina-Santiago C, Tribelli PM, Ramos JL, Molin S. 2015. Diversity of small RNAs expressed in *Pseudomonas* species. *Environ Microbiol Rep* 7:227–236. <https://doi.org/10.1111/1758-2229.12233>.
- Gomez-Lozano M, Marvig RL, Tulstrup MV, Molin S. 2014. Expression of antisense small RNAs in response to stress in *Pseudomonas aeruginosa*. *BMC Genomics* 15:783. <https://doi.org/10.1186/1471-2164-15-783>.
- Dar D, Shamir M, Mellin JR, Koutero M, Stern-Ginossar N, Cossart P, Sorek R. 2016. Term-seq reveals abundant ribo-regulation of antibiotics resistance in bacteria. *Science* 352:aad9822. <https://doi.org/10.1126/science.aad9822>.
- Poole K, Neshat S, Krebs K, Heinrichs DE. 1993. Cloning and nucleotide sequence analysis of the ferripyoverdine receptor gene *fpvA* of *Pseudomonas aeruginosa*. *J Bacteriol* 175:4597–4604. <https://doi.org/10.1128/jb.175.15.4597-4604.1993>.
- Ferrara S, Brugnoli M, De Bonis A, Righetti F, Delvillani F, Deho G, Horner D, Briani F, Bertoni G. 2012. Comparative profiling of *Pseudomonas aeruginosa* strains reveals differential expression of novel unique and conserved small RNAs. *PLoS One* 7:e36553. <https://doi.org/10.1371/journal.pone.0036553>.
- Kawano M, Reynolds AA, Miranda-Rios J, Storz G. 2005. Detection of 5'- and 3'-UTR-derived small RNAs and *cis*-encoded antisense RNAs in *Escherichia coli*. *Nucleic Acids Res* 33:1040–1050. <https://doi.org/10.1093/nar/gki256>.
- Thomason MK, Bischler T, Eisenbart SK, Forstner KU, Zhang A, Herbig A, Niesel K, Sharma CM, Storz G. 2015. Global transcriptional start site mapping using differential RNA sequencing reveals novel antisense RNAs in *Escherichia coli*. *J Bacteriol* 197:18–28. <https://doi.org/10.1128/JB.02096-14>.
- Chao Y, Vogel J. 2016. A 3' UTR-derived small RNA provides the regulatory noncoding arm of the inner membrane stress response. *Mol Cell* 61:352–363. <https://doi.org/10.1016/j.molcel.2015.12.023>.
- Guo MS, Updegrove TB, Gogol EB, Shabalina SA, Gross CA, Storz G. 2014. MicL, a new sigmaE-dependent sRNA, combats envelope stress by repressing synthesis of Lpp, the major outer membrane lipoprotein. *Genes Dev* 28:1620–1634. <https://doi.org/10.1101/gad.243485.114>.
- Cohen O, Doron S, Wurtzel O, Dar D, Edelhelt S, Karunker I, Mick E, Sorek R. 2016. Comparative transcriptomics across the prokaryotic tree of life. *Nucleic Acids Res* 44:W46–W53. <https://doi.org/10.1093/nar/gkw394>.
- Millman A, Dar D, Shamir M, Sorek R. 2017. Computational prediction of regulatory, premature transcription termination in bacteria. *Nucleic Acids Res* 45:886–893. <https://doi.org/10.1093/nar/gkw749>.
- Kalvari I, Argasinska J, Quinones-Olvera N, Nawrocki EP, Rivas E, Eddy SR, Bateman A, Finn RD, Petrov AI. 2018. Rfam 13.0: shifting to a genome-centric resource for non-coding RNA families. *Nucleic Acids Res* 46: D335–D342. <https://doi.org/10.1093/nar/gkx1038>.

36. de Kievit TR, Kakai Y, Register JK, Pesci EC, Iglewski BH. 2002. Role of the *Pseudomonas aeruginosa las* and *rhl* quorum-sensing systems in *rhlI* regulation. *FEMS Microbiol Lett* 212:101–106. [https://doi.org/10.1016/S0378-1097\(02\)00735-8](https://doi.org/10.1016/S0378-1097(02)00735-8).
37. Zuker M. 2003. Mfold web server for nucleic acid folding and hybridization prediction. *Nucleic Acids Res* 31:3406–3415. <https://doi.org/10.1093/nar/gkg595>.
38. Otaka H, Ishikawa H, Morita T, Aiba H. 2011. PolyU tail of rho-independent terminator of bacterial small RNAs is essential for Hfq action. *Proc Natl Acad Sci U S A* 108:13059–13064. <https://doi.org/10.1073/pnas.1107050108>.
39. Kery MB, Feldman M, Livny J, Tjaden B. 2014. TargetRNA2: identifying targets of small regulatory RNAs in bacteria. *Nucleic Acids Res* 42:W124–W129. <https://doi.org/10.1093/nar/gku317>.
40. Firoved AM, Boucher JC, Deretic V. 2002. Global genomic analysis of AlgU (σ^F)-dependent promoters (sigmulon) in *Pseudomonas aeruginosa* and implications for inflammatory processes in cystic fibrosis. *J Bacteriol* 184:1057–1064. <https://doi.org/10.1128/jb.184.4.1057-1064.2002>.
41. Potvin E, Sanschagrin F, Levesque RC. 2008. Sigma factors in *Pseudomonas aeruginosa*. *FEMS Microbiol Rev* 32:38–55. <https://doi.org/10.1111/j.1574-6976.2007.00092.x>.
42. Sedlyarova N, Shamovsky I, Bharati BK, Epshtein V, Chen J, Gottesman S, Schroeder R, Nudler E. 2016. sRNA-mediated control of transcription termination in *E. coli*. *Cell* 167:111–121.e113. <https://doi.org/10.1016/j.cell.2016.09.004>.
43. Morita T, Ueda M, Kubo K, Aiba H. 2015. Insights into transcription termination of Hfq-binding sRNAs of *Escherichia coli* and characterization of readthrough products. *RNA* 21:1490–1501. <https://doi.org/10.1261/ma.051870.115>.
44. Hershko-Shalev T, Odenheimer-Bergman A, Elgrably-Weiss M, Ben-Zvi T, Govindarajan S, Seri H, Papenfort K, Vogel J, Altuvia S. 2016. Gifsy-1 prophage IsrK with dual function as small and messenger RNA modulates vital bacterial machineries. *PLoS Genet* 12:e1005975. <https://doi.org/10.1371/journal.pgen.1005975>.
45. Miyakoshi M, Chao Y, Vogel J. 2015. Cross talk between ABC transporter mRNAs via a target mRNA-derived sponge of the GcvB small RNA. *EMBO J* 34:1478–1492. <https://doi.org/10.15252/emboj.201490546>.
46. Pessi G, Williams F, Hindle Z, Heurlier K, Holden MT, Camara M, Haas D, Williams P. 2001. The global posttranscriptional regulator RsmA modulates production of virulence determinants and N-acylhomoserine lactones in *Pseudomonas aeruginosa*. *J Bacteriol* 183:6676–6683. <https://doi.org/10.1128/JB.183.22.6676-6683.2001>.
47. Majdalani N, Cunniff C, Sledjeski D, Elliott T, Gottesman S. 1998. DsrA RNA regulates translation of RpoS message by an anti-antisense mechanism, independent of its action as an antisilencer of transcription. *Proc Natl Acad Sci U S A* 95:12462–12467. <https://doi.org/10.1073/pnas.95.21.12462>.
48. Majdalani N, Hernandez D, Gottesman S. 2002. Regulation and mode of action of the second small RNA activator of RpoS translation, RprA. *Mol Microbiol* 46:813–826. <https://doi.org/10.1046/j.1365-2958.2002.03203.x>.
49. Mandin P, Gottesman S. 2010. Integrating anaerobic/aerobic sensing and the general stress response through the ArcZ small RNA. *EMBO J* 29:3094–3107. <https://doi.org/10.1038/emboj.2010.179>.
50. Thomason MK, Fontaine F, De Lay N, Storz G. 2012. A small RNA that regulates motility and biofilm formation in response to changes in nutrient availability in *Escherichia coli*. *Mol Microbiol* 84:17–35. <https://doi.org/10.1111/j.1365-2958.2012.07965.x>.
51. Pearson JP, Gray KM, Passador L, Tucker KD, Eberhard A, Iglewski BH, Greenberg EP. 1994. Structure of the autoinducer required for expression of *Pseudomonas aeruginosa* virulence genes. *Proc Natl Acad Sci U S A* 91:197–201. <https://doi.org/10.1073/pnas.91.1.197>.
52. Parsek MR, Schaefer AL, Greenberg EP. 1997. Analysis of random and site-directed mutations in *rhlI*, a *Pseudomonas aeruginosa* gene encoding an acylhomoserine lactone synthase. *Mol Microbiol* 26:301–310. <https://doi.org/10.1046/j.1365-2958.1997.5741935.x>.
53. Pearson JP, Pesci EC, Iglewski BH. 1997. Roles of *Pseudomonas aeruginosa las* and *rhl* quorum-sensing systems in control of elastase and rhamnolipid biosynthesis genes. *J Bacteriol* 179:5756–5767. <https://doi.org/10.1128/jb.179.18.5756-5767.1997>.
54. Cornforth DM, Popat R, McNally L, Gurney J, Scott-Phillips TC, Ivens A, Diggle SP, Brown SP. 2014. Combinatorial quorum sensing allows bacteria to resolve their social and physical environment. *Proc Natl Acad Sci U S A* 111:4280–4284. <https://doi.org/10.1073/pnas.1319175111>.