

Opportunity and Means: Horizontal Gene Transfer from the Human Host to a Bacterial Pathogen

Mark T. Anderson and H. Steven Seifert

Department of Microbiology-Immunology, Northwestern University, Feinberg School of Medicine, Chicago, Illinois, USA

ABSTRACT The acquisition and incorporation of genetic material between nonmating species, or horizontal gene transfer (HGT), has been frequently described for phylogenetically related organisms, but far less evidence exists for HGT between highly divergent organisms. Here we report the identification and characterization of a horizontally transferred fragment of the human long interspersed nuclear element L1 to the genome of the strictly human pathogen *Neisseria gonorrhoeae*. A 685-bp sequence exhibiting 98 to 100% identity to copies of the human L1 element was identified adjacent to the *irg4* gene in some *N. gonorrhoeae* genomes. The L1 fragment was observed in ~11% of the *N. gonorrhoeae* population sampled but was not detected in *Neisseria meningitidis* or commensal *Neisseria* isolates. In addition, *N. gonorrhoeae* transcripts containing the L1 sequence were detected by reverse transcription-PCR, indicating that an L1-derived gene product may be produced. The high degree of identity between human and gonococcal L1 sequences, together with the absence of L1 sequences from related *Neisseria* species, indicates that this HGT event occurred relatively recently in evolutionary history. The identification of L1 sequences in *N. gonorrhoeae* demonstrates that HGT can occur between a mammalian host and a resident bacterium, which has important implications for the coevolution of both humans and their associated microorganisms.

IMPORTANCE The interactions between microbes and their hosts are relevant to several aspects of biology, including evolution, development, immunity, and disease. *Neisseria gonorrhoeae* serves as a particularly informative model for this interaction because it has exclusively coevolved with humans and is not known to be found in any other environment. In addition, investigation of the evolutionary relationship between *N. gonorrhoeae* and humans has practical implications, since gonorrhea is a prevalent sexually transmitted infection worldwide. This study was undertaken to characterize the horizontal transfer of genetic information from humans to *N. gonorrhoeae*, an event that has been scarcely recognized between any mammalian host and bacterial pathogen. Here we provide evidence that this genetic exchange was the result of a recent evolutionary event that has been propagated within the gonococcal population.

Received 6 January 2011 Accepted 12 January 2011 Published 15 February 2011

Citation Anderson, M. T., and H. S. Seifert. 2011. Opportunity and means: horizontal gene transfer from the human host to a bacterial pathogen. *mBio* 2(1):e00005-11. doi: 10.1128/mBio.00005-11.

Editor Stanley Maloy, San Diego State University

Copyright © 2011 Anderson and Seifert. This is an open-access article distributed under the terms of the Creative Commons Attribution-Noncommercial-Share Alike 3.0 Unported License, which permits unrestricted noncommercial use, distribution, and reproduction in any medium, provided the original author and source are credited.

Address correspondence to H. Steven Seifert, h-seifert@northwestern.edu.

Evolution of asexual organisms can occur by the accumulation and vertical transmission of mutations in an existing genetic repertoire or through the acquisition of novel genetic determinants by horizontal gene transfer (HGT). HGT has the capacity to facilitate the acquisition of complex physiological functions (1) in a single molecular event and has been estimated to account for up to 17% of the total genetic material found in *Escherichia coli* (2). HGT must therefore be regarded as an important force in the adaptation of species. HGT is particularly well known to have a diverse impact on bacterial physiology, with examples in the literature describing the horizontal acquisition of novel biosynthetic (3), virulence (1), and antibiotic resistance (4) functions by diverse bacterial species. Despite the countless interactions between commensal or pathogenic bacteria and their cognate hosts and the ever-increasing amount of available genomic sequence data, examples of bacterial integration of host genetic information are exceedingly rare. The rarity of HGT from host to bacterium may be due, in part, to potential barriers such as restriction and mod-

ification functions, the availability of genetic material, and the fitness cost of deleterious HGT events. However, such an event has the potential to impact the evolution of both the host and the microbe.

Neisseria gonorrhoeae is the causative agent of the sexually transmitted infection gonorrhea. *N. gonorrhoeae* is naturally competent for transformation and has incorporated DNA from related neisserial species (5, 6) and more divergent bacterial species (7) into its genome. Recombination of exogenous DNA occurs efficiently in *N. gonorrhoeae*, and it is postulated that this process plays an important role in the microevolution of the organism (8, 9). *N. gonorrhoeae* has a long and exclusive evolutionary history with its human host, and infection is typically localized to the urogenital tract. Therefore, sources of foreign DNA available to the gonococcus are limited to coinhabiting microorganisms or the host itself. Mammalian genomes contain numerous copies of the long interspersed nuclear element (LINE) L1. Full-length human L1 elements encode a nucleic acid binding protein and a multido-




FIG 2 PCR amplification of the nL1 fragment in *N. gonorrhoeae* isolates and nL1 transcript detection by RT-PCR. Purified genomic DNA from sequenced gonococcal isolates and human chromosomal DNA were used as templates for PCR amplification of the nL1 fragment and L1. (A) Primers IS1106for and IRGrev anneal to sequences flanking the nL1 insertion site and yield products of 1,090 and 405 bp for nL1-positive and nL1-negative alleles, respectively. Neither product was detected in DNA from strain SK-93-1035, but this isolate is not predicted to harbor the nL1 fragment. (B) Reactions using primers LINEfor and LINErev, which anneal to sequences internal to the human L1 element and the nL1 fragment. Combinations of flanking and internal primers LINErev and IS1106for (C) or LINEfor and IRGrev (D) were used to confirm the genomic location of nL1. The background bands visible in panels A and C are not due to nL1 sequence amplification, since all of the strains shown here, except PID24, DGI18, and FA6140, were also determined to be nL1 negative by DNA hybridization. Total RNA from nL1-containing isolates (DGI18, PID24, PID334, and FA6140) was used as the template in RT reaction mixtures containing (+RT) or lacking (-RT) reverse transcriptase. The resulting cDNA was amplified with internal nL1 primers L1for and L1rev. Isolates FA1090, MS11, and NCCP11945 lack the L1 fragment and were included as negative controls. Purified genomic DNA from isolate PID334 was used as a positive control. Products were obtained with cDNA generated from positive-strand (E) and negative-strand (F) RNA transcripts.

DNA during infection with *N. gonorrhoeae* versus *N. meningitidis*, or most likely an event that occurred after the two species diverged.

The absolute conservation of the nL1 sequence among the seven positive isolates is striking and suggests that this locus is under selective pressure or has not had sufficient time to degenerate. Reverse transcription (RT)-PCR using total RNA and internal L1 primers detected transcripts containing the nL1 sequence originating from both strands of DNA (Fig. 2E and F). Therefore, the HGT event that yielded nL1 has the potential to produce a novel gene product and may contribute to the observed sequence conservation. However, because the frequency of nL1 occurrence in the tested population is modest, any selective force conferred by nL1 may be small and difficult to identify experimentally. During laboratory culture, no consistent gross phenotypic differences between nL1-positive and nL1-negative strains were observed.

In addition to selective pressure, clonal expansion could also

TABLE 1 MLST profiles of the *N. gonorrhoeae* strains used in this study

Strain	MLST allele							Sequence type
	<i>abcZ</i>	<i>adk</i>	<i>aroE</i>	<i>fumC</i>	<i>gdh</i>	<i>pdhC</i>	<i>pgm</i>	
NCCP11945	109	39	170	111	148	153	65	1901
FA19	59	39	67	156	188	71	133	1892
MS11	126	39	67	78	146	153	133	6959
1291	126	39	170	158	149	531	65	8422
35/02	59	39	67	78	148	153	65	7363
FA6140	126	39	67	78	149	71	65	1927
PID1	126	39	67	156	149	71	133	8417
PID18	59	113	67	158	148	154	133	1926
PID24	109	39	170	78	228	71	65	8418
PID332	109	39	67	156	152	71	65	1594
PID334	126	39	67	156	146	71	65	8423
FA1090	109	39	67	190	147	71	65	1899
SK-92-679	59	39	67	111	149	153	65	6715
SK-93-1035	129	39	67	156	149	153	65	1595
DGI2	59	113	67	158	147	530	133	8421
DGI18	109	39	170	78	228	71	65	8418

account for the observed lack of variation of the nL1 allele among *N. gonorrhoeae* isolates. Multilocus sequencing typing (MLST) (18) was used to assess the relatedness of four nL1-positive isolates compared to 12 confirmed L1-negative isolates for which allele sequences were available. Strains DGI18 and PID24 have identical MLST profiles belonging to sequence type 8418 and may represent clonal expansion of the nL1 allele (Table 1). However, strains FA6140 and PID334 have sequence types divergent from both sequence type 8418 and each other. Since three distinct sequence types are represented among the four nL1 strains analyzed, either the presence of the nL1 fragment in these strains was the result of multiple identical HGT events or, more likely, an initial acquisition of nL1 from humans was subsequently transferred horizontally between gonococcal strains.

DISCUSSION

The high level of identity between human L1 and nL1 sequences, combined with its absence in closely related *Neisseria* species, suggests that this HGT event happened relatively recently in evolutionary history and/or that the nL1 region is under strong selective pressure. In addition, the low penetrance of nL1 within the gonococcal population is consistent with the hypothesis that nL1 was acquired recently from humans. Since numerous different L1 copies exist in any one human genome, including ones that contain sequences identical to nL1 (GenBank accession no. AC013546), analysis based on the accumulation of synonymous and nonsynonymous substitutions relative to a donor sequence is not feasible. *N. gonorrhoeae* can be found intracellularly and extracellularly associated with neutrophils and epithelial cells and thus has the opportunity to encounter host DNA during infection (19, 20). The natural competence of *N. gonorrhoeae* provides the bacterium with the means to acquire exogenous DNA, and the fact that the signature *Neisseria* DNA uptake sequence is not absolutely required for internalization of exogenous DNA (21) increases the probability of this event's occurrence. Although *N. gonorrhoeae* can invade and replicate within host cells, there is no evidence for nuclear association, making interaction with host DNA in live cells unlikely. A potential scenario in which HGT between humans and intracellular *N. gonorrhoeae* could more readily occur is after an apoptotic (22) or necrotic event in which the host DNA is

fragmented or disrupted within the dying cell. Alternatively, the active secretion of neutrophil extracellular traps (23) provides a means by which host DNA could become accessible to extracellular *N. gonorrhoeae*.

The mechanism by which this foreign genetic material was incorporated into the *N. gonorrhoeae* genome is a matter of speculation. Comparison of the insertion site between nL1-positive and nL1-negative isolates demonstrates that the recombined fragment is the result of a simple insertion with no loss or gain of flanking chromosomal information (Fig. 1). Mobilization of L1 elements within the nuclei of host cells occurs via target-primed RT (11), which requires the function of ORF1p and ORF2p and results in target site duplication. No such signature duplication is present near the nL1 site, and retrotransposition is unlikely to account for the insertion of nL1 into the *N. gonorrhoeae* genome. There is no homology between the gonococcal insertion site and the L1 sequences flanking the nL1 fragment, suggesting that direct recombination after transformation is also an unlikely mechanism for the integration of nL1. Therefore, we postulate that non-homologous end joining (NHEJ) could have occurred between a fragmented copy of L1 and the recipient region of the *N. gonorrhoeae* genome. Although *N. gonorrhoeae* lacks the NHEJ machinery possessed by some bacteria, alternative pathways of low-frequency NHEJ that have the potential to mediate HGT of exogenous sequences have been proposed (24).

Regardless of the mechanism by which HGT occurred, it is possible that nL1 has an effect on the physiology of *N. gonorrhoeae* by way of residual activity of the truncated ORF1 protein (if produced) or via polar effects of the insertion itself. More significantly, though, the identification of human sequences within the *N. gonorrhoeae* genome has important implications for modeling of the generation of genetic diversity and evolution of this and other obligate human pathogens and establishes that HGT can occur between a mammalian host and an associated microorganism.

MATERIALS AND METHODS

Bacterial strains and culture conditions. The *Neisseria* isolates used in this study and their sources are listed in Table S1 in the supplemental material. *N. gonorrhoeae* and commensal *Neisseria* species were grown on gonococcal medium base (Difco) containing Kellogg supplements as described previously (25). *N. meningitidis* strains were cultivated on gonococcal medium supplemented with 1% IsoVitalEx (Becton Dickinson). Bacteria used for the preparation of RNA were grown to exponential phase as described previously (25) in GCBL broth (1.5% proteose peptone no. 3 [Difco], 0.4% K₂HPO₄, 0.1% KH₂PO₄, 0.1% NaCl) supplemented with 1% IsoVitalEx.

Detection of the nL1 allele. All of the oligonucleotide primers used in this study are listed in Table S2 in the supplemental material. Genomic DNA was purified on QIAamp columns (Qiagen) and used for PCR and slot blot hybridizations. The nL1 genotype of the *N. gonorrhoeae* strains listed in Fig. 2 was established using the following primer combinations: IS1106for and IRGrev, LINEfor and LINErev, IS1106for and LINErev, and LINEfor and IRGrev. All other *Neisseria* isolates screened by PCR as reported in Table S1 in the supplemental material were screened by the LINEfor and LINErev primer pair at minimum. PCR products were sequenced by the Northwestern University Genomics Core Facility. A sampling of isolates of each *Neisseria* species screened was analyzed by slot blot hybridization using a 541-bp probe PCR amplified from human chromosomal DNA using primers LINEfor and LINErev. The product was labeled with digoxigenin-dUTP via the random-primed method (Roche), and hybridizations were performed using standard techniques. The binomial

distribution was used to determine the probability of observing zero nL1 events in *N. meningitidis* based on the *N. gonorrhoeae* frequency and assuming equivalent rates. The Northwestern University Biostatistics Collaboration Center assisted with the statistical analysis.

RT-PCR. Total RNA was purified from *N. gonorrhoeae* cultures using the RNeasy kit (Qiagen) and treated with RQ1 DNase (Promega) to remove genomic contamination. The LINEfor and LINErev oligonucleotides were used to prime cDNA synthesis using Superscript III reverse transcriptase (Invitrogen) according to the manufacturer's recommendations. Internal nL1 primers L1for and L1rev were used to PCR amplify cDNA products.

MLST. Seven housekeeping genes; *abcZ*, *adk*, *aroE*, *fumC*, *gdh*, *pdhC*, and *pgm*, were used to establish the sequence types of *N. gonorrhoeae* strains as described previously (18), with the exception of the oligonucleotide primers that were used for amplification and sequencing of MSLT alleles. Allele numbers were assigned by querying the *Neisseria* MLST database (18) using existing genomic sequences when available. The MLST primers listed in Table S2 in the supplemental material were used to PCR amplify and sequence (both strands) all alleles from strain PID334 and any other allele not previously established in the database. The profiles of all of the strains listed in Table 1 have been deposited in the *Neisseria* MLST database.

ACKNOWLEDGMENTS

We acknowledge M. Apicella, J. Dillard, K. Lee, R. Nicholas, P. Rice, W. Shafer, D. Trees, and W. Whittington for contributing *Neisseria* strains; J. Bodily for providing preparations of human genomic DNA; K. Clark for performing the bioinformatics screen that identified the nL1 sequence; D. Ward and M. Giovanni for facilitating the genomic sequencing of *N. gonorrhoeae* strains; M. Fitzgerald for providing genomic assemblies; M. Maiden for providing access to *N. meningitidis* genome sequences; K.-Y. Kim for assisting with the statistical analysis; and A. Chen and A. Hauser for critical reading of the manuscript.

This work was supported by NIH grants R01 AI044239 and R37 AI033493 to H.S.S. M.T.A. was partially supported by NIH fellowship F32AI080083.

SUPPLEMENTAL MATERIAL

Supplemental material for this article may be found at <http://mbio.asm.org/lookup/suppl/doi:10.1128/mBio.00005-11/-/DCSupplemental>.

Table S1, PDF file, 0.189 MB.

Table S2, PDF file, 0.042 MB.

REFERENCES

- McDaniel, T. K., and J. B. Kaper. 1997. A cloned pathogenicity island from enteropathogenic *Escherichia coli* confers the attaching and effacing phenotype on *E. coli* K-12. *Mol. Microbiol.* 23:399–407.
- Lawrence, J. G., and H. Ochman. 1997. Amelioration of bacterial genomes: rates of change and exchange. *J. Mol. Evol.* 44:383–397.
- Goldman, B. S., and R. G. Kranz. 1998. Evolution and horizontal transfer of an entire biosynthetic pathway for cytochrome c biogenesis: *Helicobacter*, *Deinococcus*, *Archae* and more. *Mol. Microbiol.* 27:871–873.
- Davies, J., and D. Davies. 2010. Origins and evolution of antibiotic resistance. *Microbiol. Mol. Biol. Rev.* 74:417–433.
- Feil, E., J. Zhou, J. Maynard Smith, and B. G. Spratt. 1996. A comparison of the nucleotide sequences of the *adk* and *recA* genes of pathogenic and commensal *Neisseria* species: evidence for extensive interspecies recombination within *adk*. *J. Mol. Evol.* 43:631–640.
- Spratt, B. G., L. D. Bowler, Q. Y. Zhang, J. Zhou, and J. M. Smith. 1992. Role of interspecies transfer of chromosomal genes in the evolution of penicillin resistance in pathogenic and commensal *Neisseria* species. *J. Mol. Evol.* 34:115–125.
- Kroll, J. S., K. E. Wilks, J. L. Farrant, and P. R. Langford. 1998. Natural genetic exchange between *Haemophilus* and *Neisseria*: intergeneric transfer of chromosomal genes between major human pathogens. *Proc. Natl. Acad. Sci. U. S. A.* 95:12381–12385.
- Fudyk, T. C., I. W. Maclean, J. N. Simonsen, E. N. Njagi, J. Kimani,

- R. C. Brunham, and F. A. Plummer. 1999. Genetic diversity and mosaicism at the *por* locus of *Neisseria gonorrhoeae*. *J. Bacteriol.* 181:5591–5599.
9. Hobbs, M. M., T. M. Alcorn, R. H. Davis, W. Fischer, J. C. Thomas, I. Martin, C. Ison, P. F. Sparling, and M. S. Cohen. 1999. Molecular typing of *Neisseria gonorrhoeae* causing repeated infections: evolution of porin during passage within a community. *J. Infect. Dis.* 179:371–381.
 10. Scott, A. F., B. J. Schmeckpeper, M. Abdelrazik, C. T. Comey, B. O'Hara, J. P. Rossiter, T. Cooley, P. Heath, K. D. Smith, and L. Margolet. 1987. Origin of the human L1 elements: proposed progenitor genes deduced from a consensus DNA sequence. *Genomics* 1:113–125.
 11. Cost, G. J., Q. Feng, A. Jacquier, and J. D. Boeke. 2002. Human L1 element target-primed reverse transcription in vitro. *EMBO J.* 21: 5899–5910.
 12. Chung, G. T., J. S. Yoo, H. B. Oh, Y. S. Lee, S. H. Cha, S. J. Kim, and C. K. Yoo. 2008. Complete genome sequence of *Neisseria gonorrhoeae* NCCP11945. *J. Bacteriol.* 190:6035–6036.
 13. Martin, S. L., and F. D. Bushman. 2001. Nucleic acid chaperone activity of the ORF1 protein from the mouse LINE-1 retrotransposon. *Mol. Cell. Biol.* 21:467–475.
 14. Khazina, E., and O. Weichenrieder. 2009. Non-LTR retrotransposons encode noncanonical RRM domains in their first open reading frame. *Proc. Natl. Acad. Sci. U. S. A.* 106:731–736.
 15. Kawai, M., I. Uchiyama, and I. Kobayashi. 2005. Genome comparison in silico in *Neisseria* suggests integration of filamentous bacteriophages by their own transposase. *DNA Res.* 12:389–401.
 16. Skaar, E. P., B. Lecuyer, A. G. Lenich, M. P. Lazio, D. Perkins-Balding, H. S. Seifert, and A. C. Karls. 2005. Analysis of the Piv recombinase-related gene family of *Neisseria gonorrhoeae*. *J. Bacteriol.* 187:1276–1286.
 17. Deitsch, K. W., J. M. Carlton, J. C. Wootton, and T. E. Wellems. 2001. Host sequences in *Plasmodium falciparum* and *Plasmodium vivax* genomic DNA: horizontal transfer or contamination artifact? *FEBS Lett.* 491: 164–165.
 18. Bennett, J. S., K. A. Jolley, P. F. Sparling, N. J. Saunders, C. A. Hart, I. M. Feavers, and M. C. Maiden. 2007. Species status of *Neisseria gonorrhoeae*: evolutionary and epidemiological inferences from multilocus sequence typing. *BMC Biol.* 5:35.
 19. Apicella, M. A., M. Ketterer, F. K. Lee, D. Zhou, P. A. Rice, and M. S. Blake. 1996. The pathogenesis of gonococcal urethritis in men: confocal and immunoelectron microscopic analysis of urethral exudates from men infected with *Neisseria gonorrhoeae*. *J. Infect. Dis.* 173:636–646.
 20. Ovcinnikov, N. M., and V. V. Delektorskij. 1971. Electron microscope studies of gonococci in the urethral secretions of patients with gonorrhoea. *Br. J. Vener. Dis.* 47:419–439.
 21. Duffin, P. M., and H. S. Seifert. 2010. DNA uptake sequence-mediated enhancement of transformation in *Neisseria gonorrhoeae* is strain dependent. *J. Bacteriol.* 192:4436–4444.
 22. Walker, P. R., and M. Sikorska. 1994. Endonuclease activities, chromatin structure, and DNA degradation in apoptosis. *Biochem. Cell Biol.* 72: 615–623.
 23. Brinkmann, V., U. Reichard, C. Goosmann, B. Fauler, Y. Uhlemann, D. S. Weiss, Y. Weinrauch, and A. Zychlinsky. 2004. Neutrophil extracellular traps kill bacteria. *Science* 303:1532–1535.
 24. Chayot, R., B. Montagne, D. Mazel, and M. Ricchetti. 2010. An end-joining repair mechanism in *Escherichia coli*. *Proc. Natl. Acad. Sci. U. S. A.* 107:2141–2146.
 25. Stohl, E. A., A. K. Criss, and H. S. Seifert. 2005. The transcriptome response of *Neisseria gonorrhoeae* to hydrogen peroxide reveals genes with previously uncharacterized roles in oxidative damage protection. *Mol. Microbiol.* 58:520–532.