



# The Transferable Resistome of Produce

Khald Blau,<sup>a</sup> Antje Bettermann,<sup>a</sup> Sven Jechalke,<sup>b</sup> Eva Fornefeld,<sup>a</sup> Yann Vanrobaeys,<sup>c,d</sup> Thibault Stalder,<sup>c,d</sup> Eva M. Top,<sup>c,d</sup> Kornelia Smalla<sup>a</sup>

<sup>a</sup>Julius Kühn-Institut, Federal Research Centre for Cultivated Plants, Braunschweig, Germany

<sup>b</sup>Justus Liebig University Giessen, Institute for Phytopathology, Gießen, Germany

<sup>c</sup>Department of Biological Sciences, University of Idaho, Moscow, Idaho, USA

<sup>d</sup>Institute for Bioinformatics and Evolutionary Studies, University of Idaho, Moscow, Idaho, USA

**ABSTRACT** Produce is increasingly recognized as a reservoir of human pathogens and transferable antibiotic resistance genes. This study aimed to explore methods to characterize the transferable resistome of bacteria associated with produce. Mixed salad, arugula, and cilantro purchased from supermarkets in Germany were analyzed by means of cultivation- and DNA-based methods. Before and after a nonselective enrichment step, tetracycline (TET)-resistant *Escherichia coli* were isolated and plasmids conferring TET resistance were captured by exogenous plasmid isolation. TET-resistant *E. coli* isolates, transconjugants, and total community DNA (TC-DNA) from the microbial fraction detached from leaves or after enrichment were analyzed for the presence of resistance genes, class 1 integrons, and various plasmids by real-time PCR and PCR-Southern blot hybridization. Real-time PCR primers were developed for IncI and IncF plasmids. TET-resistant *E. coli* isolated from arugula and cilantro carried IncF, IncI1, IncN, IncHI1, IncU, and IncX1 plasmids. Three isolates from cilantro were positive for IncN plasmids and *bla*<sub>CTX-M-1</sub>. From mixed salad and cilantro, IncF, IncI1, and IncP-1 $\beta$  plasmids were captured exogenously. Importantly, whereas direct detection of IncI and IncF plasmids in TC-DNA failed, these plasmids became detectable in DNA extracted from enrichment cultures. This confirms that cultivation-independent DNA-based methods are not always sufficiently sensitive to detect the transferable resistome in the rare microbiome. In summary, this study showed that an impressive diversity of self-transmissible multiple resistance plasmids was detected in bacteria associated with produce that is consumed raw, and exogenous capturing into *E. coli* suggests that they could transfer to gut bacteria as well.

**IMPORTANCE** Produce is one of the most popular food commodities. Unfortunately, leafy greens can be a reservoir of transferable antibiotic resistance genes. We found that IncF and IncI plasmids were the most prevalent plasmid types in *E. coli* isolates from produce. This study highlights the importance of the rare microbiome associated with produce as a source of antibiotic resistance genes that might escape cultivation-independent detection, yet may be transferred to human pathogens or commensals.

**KEYWORDS** *Escherichia coli*, IncF, IncI, antibiotic resistance, horizontal gene transfer, real-time PCR

Despite its benefit to human health, consumption of produce is increasingly recognized as a source of pathogenic bacteria, antibiotic-resistant bacteria (ARB), and antibiotic resistance genes (ARGs) associated with mobile genetic elements (MGEs) (1–5). Recently, several foodborne disease outbreaks have been associated with produce contamination worldwide (5–9). The microbiome of produce is important for plant health and vigor and was shown to be highly dynamic during growth and postharvest (10), but can also contain potentially pathogenic bacteria from human and animal


Received 14 June 2018 Accepted 20 September 2018 Published 6 November 2018

**Citation** Blau K, Bettermann A, Jechalke S, Fornefeld E, Vanrobaeys Y, Stalder T, Top EM, Smalla K. 2018. The transferable resistome of produce. *mBio* 9:e01300-18. <https://doi.org/10.1128/mBio.01300-18>.

**Editor** Mark J. Bailey, CEH-Oxford

**Copyright** © 2018 Blau et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Kornelia Smalla, [kornelia.smalla@julius-kuehn.de](mailto:kornelia.smalla@julius-kuehn.de).



**FIG 1** Flow diagram of the experimental setup of this study to evaluate the use of culture-dependent and -independent approaches to characterize the transferable resistome of bacteria associated with produce.

sources, including *Escherichia coli* strains (11). Contamination can occur preharvest (i.e., through organic fertilizers, soil, wild animals, or contaminated irrigation water) and postharvest (12, 13).

Antibiotic resistance in bacterial pathogens has increased globally due to the widespread use and misuse of antibiotics (14–17). Antibiotic resistance levels in *E. coli* are useful indicators of overall resistance levels of bacteria on foods and in animals and humans (11). Antibiotic resistance and ARGs have been documented for enteric bacteria from various types of produce, which could facilitate the dissemination of resistant bacteria to a wider community of people (1, 2, 4, 16, 18, 19). If ARGs are localized on MGEs such as plasmids or conjugative transposons they can be transferred horizontally to pathogens (20). Horizontal gene transfer (HGT) takes place at sites with high cell densities of plasmid donors and recipients, nutrient availability, and selective pressure. The phytosphere, including the rhizosphere and the phyllosphere, have been reported as hot spots of HGT (21). The plasmid-mediated resistome of produce bacteria might provide the enterobacteria with ARGs in the intestine under selective conditions. Conjugative plasmids can often confer resistance not only toward multiple antibiotics but also toward heavy metal compounds or disinfectants, making coselection possible (22–25). Although plasmids belonging to the incompatibility groups IncF and IncI have a narrow host range (NHR), they are assumed to be important for the dissemination of ARGs in *E. coli* and other *Enterobacteriaceae* (26, 27). Most importantly, resistance- and virulence-associated traits of *E. coli* isolates were almost exclusively found on IncF group plasmids (28–30). However, no real-time PCR (RT-PCR) systems that allow the cultivation-independent detection and quantification of these plasmids in total community DNA (TC-DNA) are available.

In this study, culture-dependent and -independent approaches were employed to assess the transferable resistome of bacteria associated with produce (Fig. 1). We focused on tetracycline (TET) resistance because of the large amounts of tetracyclines used in animal husbandries resulting in a high load released into the agro-ecosystem via organic fertilizers (31). TET-resistant *E. coli* was isolated from produce directly after purchase and after seven days of storage by selective plating with and without prior

nonselective enrichment. In addition, transferable TET resistance plasmids were captured into *E. coli* recipient strains using the so-called exogenous plasmid isolation method (32). New real-time PCR primers were developed for the detection and quantification of IncF and IncI plasmids. TC-DNA was also extracted from the microbial fraction detached from produce or after nonselective enrichment to detect and quantify the abundance of ARGs and MGEs.

## RESULTS

**Phenotypic and genotypic characterization of TET-resistant *E. coli* isolates.** To find out whether produce was a source of antibiotic-resistant *E. coli*, we determined the occurrence and resistance profiles of TET-resistant *E. coli* isolated from 24 samples of produce directly or after an overnight enrichment step. The phenotypic characterization of a total of 63 TET-resistant *E. coli* isolates from cilantro ( $n = 54$ ), arugula ( $n = 7$ ), and mixed salad ( $n = 2$ ), of which 50 were recovered after nonselective enrichment and 13 without enrichment (20.6%) revealed an impressive diversity (Table 1).

Almost all *E. coli* isolates were resistant to antibiotics from at least one class, and two isolates were resistant to eight antibiotic classes, tetracyclines (TET and D), penicillins (AM and AMX), third generation cephalosporins (CTX and CRO), fluoroquinolones (CIP, OFX, and NA), aminoglycosides (GM and S), sulfonamides (SD), phenicols (C), and trimethoprim (TMP). Most of the TET-resistant *E. coli* also displayed resistance to ampicillin and amoxicillin (84%) and trimethoprim (73%). Resistances to ofloxacin, ciprofloxacin, sulfadiazine, and streptomycin were also common. We tested all of the isolates for the production of extended-spectrum beta-lactamases (ESBLs) with the double-disc diffusion test (DDT) and found three ESBL-producing *E. coli* which were isolated from two of the cilantro samples.

We then genotypically characterized the collection of *E. coli* isolates for the presence of various resistance genes [*tet(A)*, *strA*, *sul1*, *sul2*, *sul3*, *aadA*, *qacE* and/or *qacEΔ1* (*qacE/qacEΔ1*), *merRTΔP*, *bla* genes (TEM, CTX-M, and SHV), *qnr* genes (*qnrA*, *qnrB*, and *qnrS*)] and integrase genes *int1* and *int2* by RT-PCR or regular PCR of genomic DNA (Table 1). The most commonly detected ARG was the tetracycline resistance gene *tet(A)*, which was found in 59 out of 63 isolates. A total of 10 isolates were positive for the sulfonamide resistance genes *sul1*, 14 for *sul2*, and five for *sul3*. The combinations of *sul1* and *sul2*, *sul2* and *sul3*, and *sul1* and *sul3* were detected in seven, three, and one isolate, respectively. All three *sul* genes were found in one TET-resistant *E. coli* isolate from cilantro. The *qnrB* and *qnrS* genes encoding fluoroquinolone resistance were detected alone or in combination in one and 38 isolates, respectively. The *bla*<sub>TEM</sub> genes encoding resistance to ampicillin and amoxicillin were detected in 82.5% of TET-resistant *E. coli* isolates. The *bla*<sub>CTX-M-1</sub> gene encoding ESBL resistance was detected in only three isolates and was found in combination with *bla*<sub>TEM</sub> genes in two *E. coli* from cilantro. The *bla*<sub>SHV</sub> gene encoding ESBL resistance was not detected in any of the isolates. For the streptomycin/spectinomycin resistance genes, *aadA* (24 isolates) was most common, followed by *strA* (21 isolates) and *aadA* and *strA* (three isolates).

The class 1 integron integrase gene *int1* was detected in 50 isolates, while the class 2 integron integrase gene *int2* was not detected at all. Although *qacEΔ1* encoding quaternary ammonium compound resistance is a typical component of class 1 integrons, the *qacE* and/or *qacEΔ1* genes were detected in only 23 isolates, suggesting a large proportion of atypical class 1 integrons. Interestingly, *merRTΔP* encoding regulation, transport, and extracellular mercury-binding was detected in 12 isolates. These findings show that produce can be a source of multidrug-resistant *E. coli* isolates.

**Characterization of plasmids in TET-resistant *E. coli* isolates.** To test if the TET-resistant *E. coli* isolates recovered from produce harbor plasmids and to assign them to known plasmid groups, their genomic DNA was screened by TaqMan probe-based RT-PCR systems for IncF and IncI plasmids and by PCR-based replicon typing (PBRT) (Table 1). All isolates that were positive by RT-PCR targeting the IncF (*traI* gene) were also identified by replicon typing as IncF, confirming the specificity of the novel TaqMan RT-PCR system. However, PBRT also allowed assignment to the different IncF

**TABLE 1** Characterization of representative tetracycline-resistant *E. coli* isolates from produce

<i>E. coli</i> isolates <sup>a</sup>	Sample source <sup>b</sup>	Time point (day)	Inc group <sup>c</sup>	<i>bla</i> genes	Resistance and integrase genes	Antibiotic resistance profile <sup>d</sup>
EK2.1 <sup>D</sup>	Ci	0	FII <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>tet(A)</i> , <i>qnrS</i>	AM, AMX, TET, CIP, OFX
EK2.2 <sup>E</sup>	Ci	0	U <sup>h</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>sul2</i> , <i>strA</i> , <i>qnrS</i>	AM, AMX, TET, S, TMP, SD, CIP, OFX
EK2.5 <sup>D</sup>	Ci	0	I1 <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>tet(A)</i> , <i>qnrS</i>	AM, AMX, TET, CIP
EK2.7 <sup>D</sup>	Ci	0	U <sup>h</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>sul2</i> , <i>strA</i> , <i>qnrS</i>	AM, AMX, TET, S, TMP, SD, OFX
EK2.8 <sup>D</sup>	Ci	0	I1 <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>sul2</i> , <i>strA</i>	AM, AMX, TET, S, TMP
EK2.11 <sup>D</sup>	Ci	0	I1 <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>tet(A)</i>	AM, AMX, TET
EK2.15 <sup>D</sup>	Ci	0	I1 <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>tet(A)</i> , <i>sul2</i>	AM, AMX, TET
EK2.16 <sup>D</sup>	Ci	0	N <sup>g</sup>	<i>bla</i> <sub>TEM</sub>	<i>tet(A)</i> , <i>sul2</i> , <i>sul3</i> , <i>strA</i>	AM, AMX, TET, S
EK2.18 <sup>D</sup>	Ci	0	X1 <sup>h</sup>	<i>bla</i> <sub>TEM</sub>	<i>tet(A)</i> , <i>sul2</i> , <i>sul3</i> , <i>strA</i>	AM, AMX, TET, S
EK2.19 <sup>E</sup>	Ci	0	FII <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>merRTΔP</i> , <i>sul2</i> , <i>aadA</i>	AM, AMX, TET, S, TMP, D, GM, KM
EK2.20 <sup>E</sup>	Ci	0	HI1 <sup>h</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>sul2</i> , <i>sul3</i> , <i>aadA</i> , <i>qnrS</i>	AM, AMX, TET, S, TMP, SD, GM, OFX, C
EK2.21 <sup>E</sup>	Ci	0	X1 <sup>h</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i>	AM, AMX, TET, OFX
EK2.22 <sup>E</sup>	Ci	0	X1 <sup>h</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>strA</i> , <i>qnrS</i>	AM, AMX, TET, S, TMP, SD, OFX
EK2.25 <sup>E</sup>	Ci	0	U <sup>h</sup> , X1 <sup>h</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>sul2</i> , <i>strA</i> , <i>qnrS</i>	AM, AMX, TET, S, TMP, SD, CIP, OFX
EK2.26 <sup>E</sup>	Ci	0	U <sup>h</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>sul1</i> , <i>aadA</i> , <i>qacE/qacEΔ1</i> , <i>strA</i>	AM, AMX, TET, S, TMP, SD, CIP, NA, OFX
EK2.29 <sup>E,k</sup>	Ci	0	N <sup>g</sup>	<i>bla</i> <sub>TEM</sub> , <i>bla</i> <sub>CTX-M-1</sub>	<i>int11</i> , <i>tet(A)</i> , <i>sul1</i> , <i>strA</i> , <i>qnrS</i>	AM, AMX, TET, S, TMP, SD, CRO, CTX, OFX, CIP
EK2.30 <sup>E</sup>	Ci	0	U <sup>h</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>sul1</i> , <i>sul3</i> , <i>aadA</i> , <i>qnrS</i>	AM, AMX, TET, TMP, SD, C
EK3.33 <sup>E</sup>	Ci	0	X1 <sup>h</sup>		<i>int11</i> , <i>sul1</i> , <i>strA</i>	TET, D
EK3.34 <sup>D</sup>	Ci	0	FII <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>sul1</i> , <i>strA</i>	AM, AMX, TET, S, TMP, SD, NA, OFX
EK3.35 <sup>D</sup>	Ci	0	FII <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>tet(A)</i> , <i>sul1</i>	AM, AMX, TET, S, TMP, SD, NA
EK3.36 <sup>E</sup>	Ci	0	FIB <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>merRTΔP</i> , <i>sul1</i> , <i>aadA</i>	AM, AMX, TET, S, TMP, D, GM, KM
EK3.43 <sup>E,k</sup>	Ci	0	N <sup>g</sup>	<i>bla</i> <sub>CTX-M-1</sub>	<i>tet(A)</i> , <i>sul1</i> , <i>qnrS</i>	AM, AMX, TET, S, D, GM, CTX, OFX
EK3.44 <sup>E,k</sup>	Ci	7	N <sup>g</sup>	<i>bla</i> <sub>TEM</sub> , <i>bla</i> <sub>CTX-M-1</sub>	<i>int11</i> , <i>tet(A)</i> , <i>merRTΔP</i> , <i>sul1</i> , <i>strA</i> , <i>qnrS</i>	AM, AMX, TET, S, TMP, SD, CRO, CTX, C, NA, CIP, OFX
EK5.16 <sup>E</sup>	Ci	7	FII <sup>f</sup> , I1 <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>aadA</i> , <i>qacE/qacEΔ1</i> , <i>qnrS</i>	AM, AMX, TET, S, TMP, D, OFX
EK5.19 <sup>E</sup>	Ci	7	X1 <sup>h</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>sul2</i> , <i>strA</i> , <i>qnrS</i>	AM, AMX, TET, S, TMP, SD, D, C
EK5.20 <sup>E</sup>	Ci	7	FII <sup>f</sup> , I1 <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>aadA</i> , <i>qacE/qacEΔ1</i> , <i>qnrS</i>	AM, AMX, TET, S, TMP, OFX
EK5.25 <sup>E</sup>	Ci	7	FII <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>qacE/qacEΔ1</i> , <i>qnrS</i>	AM, AMX, TET, TMP, OFX
EK5.28 <sup>E</sup>	Ci	7	FII <sup>f</sup> , I1 <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>aadA</i> , <i>qacE/qacEΔ1</i> , <i>qnrS</i>	AM, AMX, TET, S, TMP, CIP, OFX
EK5.30 <sup>E</sup>	Ci	7	FII <sup>f</sup> , FIB <sup>f</sup> , I1 <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>sul1</i> , <i>aadA</i> , <i>qacE/qacEΔ1</i> , <i>qnrS</i>	AM, AMX, TET, TMP, SD, D, CIP, NA, OFX
EK5.32 <sup>E</sup>	Ci	7	X1 <sup>h</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>sul1</i> , <i>sul2</i> , <i>strA</i> , <i>qnrS</i>	AM, AMX, TET, S, TMP, SD, OFX, C
EK5.40 <sup>E</sup>	Ci	7	X1 <sup>h</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>sul2</i> , <i>strA</i> , <i>qnrS</i>	AM, AMX, TET, S, TMP, SD, C
EK7.6 <sup>E</sup>	Ci	7	U <sup>h</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i>	AM, AMX, TET, TMP, D
EK7.7 <sup>E</sup>	Ci	7	I1 <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>qacE/qacEΔ1</i> , <i>aadA</i>	AM, AMX, TET, TMP
EK7.9 <sup>E</sup>	Ci	7	X1 <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>merRTΔP</i> , <i>sul2</i> , <i>qnrS</i>	AM, AMX, TET, SD, CRO, D, CIP, C, GM, OFX
EK7.10 <sup>E</sup>	Ci	7	HI1 <sup>h</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>merRTΔP</i> , <i>sul2</i> , <i>qnrS</i>	AM, AMX, TET, SD, CRO, D, CIP, C, GM, OFX
EK7.11 <sup>E</sup>	Ci	7	FII <sup>f</sup> , I1 <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>aadA</i> , <i>qacE/qacEΔ1</i> , <i>qnrS</i>	AM, AMX, TET, TMP, CIP
EK7.12 <sup>E</sup>	Ci	7	FII <sup>f</sup> , I1 <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>aadA</i> , <i>qacE/qacEΔ1</i> , <i>qnrS</i>	AM, AMX, TET, TMP
EK7.13 <sup>E</sup>	Ci	7	FIB <sup>f</sup> , I1 <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>aadA</i> , <i>qacE/qacEΔ1</i> , <i>qnrS</i>	AM, AMX, TET, TMP, OFX
EK7.14 <sup>E</sup>	Ci	7	FII <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>aadA</i> , <i>qacE/qacEΔ1</i> , <i>qnrS</i>	AM, AMX, TET, TMP
EK7.15 <sup>E</sup>	Ci	7	FII <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>sul1</i> , <i>sul2</i> , <i>aadA</i> , <i>qacE/qacEΔ1</i> , <i>qnrS</i>	AM, AMX, TET, S, TMP, SD, CIP, GM
EK7.16 <sup>E</sup>	Ci	7	FII <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>sul1</i> , <i>sul2</i> , <i>strA</i> , <i>qacE/qacEΔ1</i> , <i>qnrS</i>	AM, AMX, TET, S, TMP, SD, CIP, OFX
EK7.17 <sup>E</sup>	Ci	7	FII <sup>f</sup> , X1 <sup>h</sup>	<i>bla</i> <sub>TEM</sub>	<i>tet(A)</i> , <i>qnrS</i>	AM, AMX, TET, CIP, OFX
EK7.18 <sup>E</sup>	Ci	7	FII <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>sul2</i> , <i>strA</i> , <i>qnrS</i>	AM, AMX, TET, S, TMP, SD, CIP, OFX
EK7.19 <sup>E</sup>	Ci	7	FIB <sup>f</sup> , I1 <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>aadA</i> , <i>qacE/qacEΔ1</i> , <i>qnrS</i>	AM, AMX, TET, TMP
EK7.21 <sup>E</sup>	Ci	7	FIB <sup>f</sup> , I1 <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>aadA</i> , <i>qacE/qacEΔ1</i> , <i>qnrS</i>	AM, AMX, TET, TMP, OFX
EK7.22 <sup>E</sup>	Ci	7	FII <sup>f</sup> , FIB <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>sul1</i> , <i>sul2</i> , <i>sul3</i> , <i>aadA</i> , <i>strA</i> , <i>qacE/qacEΔ1</i> , <i>qnrS</i>	AM, AMX, TET, S, TMP, SD, CIP, GM, NA, OFX
EK7.24 <sup>E</sup>	Ci	7	FII <sup>f</sup> , I1 <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>sul1</i> , <i>sul2</i> , <i>aadA</i> , <i>strA</i> , <i>qacE/qacEΔ1</i> , <i>qnrB</i> , <i>qnrS</i>	AM, AMX, TET, S, TMP, SD, CIP, OFX, C
EK7.26 <sup>E</sup>	Ci	7	FIB <sup>f</sup> , I1 <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>aadA</i> , <i>qacE/qacEΔ1</i> , <i>qnrS</i>	AM, AMX, TET, TMP, CIP, OFX
EK7.27 <sup>E</sup>	Ci	7	X1 <sup>h</sup>	<i>bla</i> <sub>TEM</sub>	<i>tet(A)</i> , <i>merRTΔP</i> , <i>sul2</i> , <i>qnrS</i>	AM, AMX, TET, SD, CRO, D, CIP, C, GM, OFX
EK7.28 <sup>E</sup>	Ci	7	FIB <sup>f</sup> , I1 <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>aadA</i> , <i>qacE/qacEΔ1</i> , <i>qnrS</i>	AM, AMX, TET, TMP
EK7.29 <sup>E</sup>	Ci	7	FII <sup>f</sup> , FIB <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>sul1</i> , <i>aadA</i> , <i>qacE/qacEΔ1</i> , <i>qnrS</i>	AM, AMX, TET, S, TMP, SD, D, CIP, NA, OFX
EK7.30 <sup>E</sup>	Ci	7	FII <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>sul1</i> , <i>sul2</i> , <i>aadA</i> , <i>qacE/qacEΔ1</i> , <i>qnrS</i>	AM, AMX, TET, S, TMP, SD, GM
EK6.1 <sup>E</sup>	Ci	7	FII <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>sul2</i> , <i>strA</i>	TET, S
EK6.3 <sup>E</sup>	Ci	7	FIB <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>tet(A)</i> , <i>sul2</i> , <i>strA</i>	TET, S
M1 <sup>E</sup>	MS	0	ND	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>sul1</i> , <i>sul2</i> , <i>qacE/qacEΔ1</i> , <i>strA</i> , <i>merRTΔP</i>	AM, AMX, TET, S, TMP, SD, D
M19 <sup>E</sup>	MS	0	ND	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>sul1</i> , <i>sul2</i> , <i>qacE/qacEΔ1</i> , <i>strA</i>	AM, AMX, TET, S, TMP, SD, D
RE1 <sup>D</sup>	A	0	FII <sup>f</sup>			TET, D
RE4 <sup>D</sup>	A	0	I1 <sup>f</sup>		<i>int11</i> , <i>tet(A)</i> , <i>sul3</i> , <i>aadA</i> , <i>merRTΔP</i>	TET, S, TMP, SD, D, CIP, NA
RE9 <sup>D</sup>	A	0	I1 <sup>f</sup>		<i>int11</i> , <i>tet(A)</i> , <i>sul3</i> , <i>aadA</i> , <i>merRTΔP</i>	TET, S, TMP, SD, D, CIP, NA
RE10 <sup>D</sup>	A	0	I1 <sup>f</sup>		<i>int11</i> , <i>tet(A)</i> , <i>sul3</i> , <i>aadA</i> , <i>merRTΔP</i>	TET, S, TMP, SD, D, CIP, NA
RE11 <sup>E</sup>	A	0	I1 <sup>f</sup>		<i>int11</i> , <i>tet(A)</i> , <i>sul3</i> , <i>aadA</i> , <i>merRTΔP</i>	TET, S, TMP, SD, D, CIP, NA
RE14 <sup>E</sup>	A	7	FII <sup>f</sup>			TET, D
RE19 <sup>E</sup>	A	7	I1 <sup>f</sup>		<i>int11</i> , <i>tet(A)</i> , <i>sul3</i> , <i>aadA</i> , <i>merRTΔP</i>	TET, S, TMP, SD, D, CIP, NA

<sup>a</sup>D, direct plating; E, enrichment.<sup>b</sup>Ci, cilantro; MS, mixed salad; A, arugula.<sup>c</sup>l, detected by RT-PCR and PBRT; f, detected by RT-PCR; g, detected by PCR; h, detected by PBRT; k, conjugal transfer into *E. coli* CV601; ND, not detected.<sup>d</sup>AM, ampicillin; AMX, amoxicillin; TET, tetracycline; CIP, ciprofloxacin; OFX, ofloxacin; S, streptomycin; TMP, trimethoprim; SD, sulfadiazine; D, doxycycline; GM, gentamicin; KM, kanamycin; C, chloramphenicol; NA, nalidixic acid; CRO, ceftriaxone; CTX, cefotaxime.

subgroups. Furthermore, other plasmids were also identified by PBRT or RT-PCR (*korB*, specific for IncP-1 plasmids) or PCR (IncN). A summary of the plasmid/replicon types detected among the 63 representative TET-resistant *E. coli* isolates is given in Table 1. For cilantro and arugula, almost all TET-resistant *E. coli* isolates contained plasmids (61 out of 63), but the plasmids detected in the two isolates from mixed salad could not be assigned using RT-PCR or PBRT. In most isolates ( $n = 45$ ), one plasmid type was detected, but some had two ( $n = 15$ ) or three ( $n = 1$ ) plasmids. Plasmids from seven different Inc groups were found in the 63 *E. coli* isolates, IncFII ( $n = 21$ ), IncI1 ( $n = 17$ ), IncX1 ( $n = 11$ ), IncFIB ( $n = 10$ ), IncU ( $n = 6$ ), IncN ( $n = 4$ ), and IncHI1 ( $n = 2$ ). All Inc groups were found in *E. coli* isolates from cilantro, whereas only two Inc groups were found in isolates from arugula, IncI1 ( $n = 5$ ) and IncFII ( $n = 2$ ). Plasmids of the IncF groups (FII and FIB) were the predominant types, followed by IncI1 and IncX1 plasmids. The combination of replicon types IncFII and IncFIB was detected in two isolates, whereas the combination of replicon types IncFII and IncI1 and the combination of IncFIB and IncI1 were found in six and five isolates, respectively. In one isolate from cilantro, the combination IncFII, IncFIB and IncI1 was detected. IncI2 plasmids were not detected in any of the *E. coli* isolates.

**Conjugal transfer of antibiotic resistance.** Conjugation experiments were conducted in order to determine the potential transfer of antibiotic resistances to other bacteria. Conjugal transfer experiments were performed using TET-resistant *E. coli* isolates positive for ESBL (EK2.29, EK3.43, and EK3.44) as donors and kanamycin- and rifampin-resistant *E. coli* CV601 as a recipient at 37°C. We selected transconjugants on LB plates containing tetracycline and cefotaxime, which corresponded to phenotypes of the donors. The transfer of the resistance phenotypes was successful.

**Phenotypic and genotypic characterization of plasmids captured via exogenous isolation.** We further investigated the presence of transferable plasmids in produce by capturing TET resistance plasmids from nonselective enrichment cultures of fresh leaves from cilantro, mixed salad, or arugula by exogenous plasmid isolation into *E. coli* CV601. TET-resistant transconjugants were captured only on day 0 but not on day 7. The transfer frequencies of TET-resistant transconjugants were  $1.73 \times 10^{-7}$ ,  $1.55 \times 10^{-4}$ , and  $4.66 \times 10^{-9}$  per recipient in cilantro, mixed salad, and arugula, respectively. While all transconjugants obtained from cilantro ( $n = 27$ ) and arugula ( $n = 23$ ) were characterized, a total of only 41 transconjugants from mixed salad was analyzed due to the high number of transconjugants obtained. Based on initial phenotypic and genotypic analyses, 15 representative out of 91 TET-resistant transconjugants from produce (cilantro,  $n = 12$ ; arugula,  $n = 1$ ; mixed salad,  $n = 2$ ) were selected for further characterization. The majority of these transconjugants acquired resistance to at least two antibiotic classes, and all were resistant to tetracycline, ampicillin, and amoxicillin. The *bla*<sub>TEM</sub> genes encoding ampicillin and amoxicillin resistances were detected in 86.7% of TET-resistant transconjugants (Table 2). The tetracycline resistance gene *tet(A)* was found in 13 out of 15 transconjugants from cilantro and arugula but not from mixed salad, while *tet(Q)* was detected in only one plasmid (pBMS1) isolated from the mixed salad. Four tetracycline resistance plasmids (pBC1.1, pBC1.3, pBC1.9, and pBC1.12) captured from cilantro carried the insertion sequence IS1071, class 1 integrons (*int1*) and tetracycline resistance gene *tet(A)*, but also encoded resistance to ampicillin (*bla*<sub>TEM</sub>), and mercury compounds (*merRTΔP*). Eight plasmids from cilantro transconjugants (pBC2.1, pBC2.2, pBC2.3, pBC2.4, pBC2.6, pBC2.8, pBC2.11, and pBC2.15) carried *tet(A)*, *qnrS*, and *bla*<sub>TEM</sub> and two of the plasmids (pBC2.1 and pBC2.4) carried in addition *sul1* and *sul2*, respectively. Two TET resistance plasmids (pBMS1 and pBMS4) captured from mixed salad carried *sul1*, *strA*, *merRTΔP*, *bla*<sub>TEM</sub>, and *int1*. One plasmid (pBA1) captured from arugula carried *bla*<sub>TEM</sub> and *tet(A)* (Table 2). Thus, this approach demonstrates that transferable multidrug resistance plasmids were easily captured by *E. coli* CV601, a process that might also occur in the human gut.

**Identification of exogenously isolated plasmids.** The newly developed TaqMan probe-based RT-PCR assay was used to screen the TET-resistant transconjugants for the

**TABLE 2** Characterization of representative tetracycline resistant *E. coli* CV601 transconjugants captured from produce

TET <sup>r</sup> <i>E. coli</i> CV601 transconjugants <sup>a</sup>	Sample source <sup>b</sup>	Inc groups <sup>c</sup>	<i>bla</i> genes	Resistance, integrase genes and IS <sup>d</sup>	Antibiotic resistance profile <sup>e</sup>
pBC1.1	Ci	P-1β <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>merRTΔP</i> , <i>qacE/qacEΔ1</i> , IS1071	TET, AM, AMX, D
pBC1.3	Ci	P-1β <sup>f</sup>		<i>int11</i> , <i>tet(A)</i> , <i>merRTΔP</i> , <i>qacE/qacEΔ1</i> , IS1071	TET, AM, AMX, D
pBC1.9	Ci	P-1β <sup>f</sup> , FII <sup>l</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(A)</i> , <i>merRTΔP</i> , <i>qacE/qacEΔ1</i> , IS1071	TET, AM, AMX, D
pBC1.12	Ci	P-1β <sup>f</sup> , FII <sup>l</sup>		<i>int11</i> , <i>tet(A)</i> , <i>merRTΔP</i> , <i>strA</i> , <i>qacE/qacEΔ1</i> , IS1071	TET, AM, AMX, D, S
pBC2.1	Ci	FIB <sup>l</sup>	<i>bla</i> <sub>TEM</sub>	<i>tet(A)</i> , <i>sul1</i> , <i>qnrS</i>	TET, AM, AMX, D, CIP, NA, OFX, C
pBC2.2	Ci	FIB <sup>l</sup>	<i>bla</i> <sub>TEM</sub>	<i>tet(A)</i> , <i>qnrS</i>	TET, AM, AMX, D, CIP, OFX
pBC2.3	Ci	FIB <sup>l</sup> , I1 <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>tet(A)</i> , <i>qnrS</i>	TET, AM, AMX, D, CIP, OFX
pBC2.4	Ci	FIB <sup>l</sup>	<i>bla</i> <sub>TEM</sub>	<i>tet(A)</i> , <i>sul2</i> , <i>qnrS</i>	TET, AM, AMX, D, CIP, NA, OFX
pBC2.6	Ci	FIB <sup>l</sup>	<i>bla</i> <sub>TEM</sub>	<i>tet(A)</i> , <i>qnrS</i>	TET, AM, AMX, D, CIP, OFX, C
pBC2.8	Ci	FIB <sup>l</sup>	<i>bla</i> <sub>TEM</sub>	<i>tet(A)</i> , <i>qnrS</i>	TET, AM, AMX, D, CIP, NA, OFX
pBC2.11	Ci	FII <sup>l</sup>	<i>bla</i> <sub>TEM</sub>	<i>tet(A)</i> , <i>qnrS</i>	TET, AM, AMX, D, CIP, OFX
pBC2.15	Ci	I1 <sup>f</sup>	<i>bla</i> <sub>TEM</sub>	<i>tet(A)</i> , <i>qnrS</i>	TET, AM, AMX, D, CIP, OFX
pBMS1	MS	FII <sup>l</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>tet(Q)</i> , <i>sul1</i> , <i>strA</i> , <i>merRTΔP</i>	TET, AM, AMX, D, TMP, C, S, SD
pBMS4	MS	FII <sup>l</sup>	<i>bla</i> <sub>TEM</sub>	<i>int11</i> , <i>sul1</i> , <i>strA</i> , <i>merRTΔP</i> ,	TET, AM, AMX, D, TMP, C, S, SD
pBA1	A	ND	<i>bla</i> <sub>TEM</sub>	<i>tet(A)</i>	TET, AM, AMX, D, TMP, C, CIP

*E. coli* CV601 (recipient)

<sup>a</sup>Superscript r indicates resistance to the antibiotic.

<sup>b</sup>Ci, cilantro; MS, mixed salad; A, arugula.

<sup>c</sup>l, detected by RT-PCR and PBRT; f, detected by RT-PCR; ND, not detected.

<sup>d</sup>IS, insertion sequence.

<sup>e</sup>TET, tetracycline; AM, ampicillin; AMX, amoxicillin; D, doxycycline; S, streptomycin; CIP, ciprofloxacin; NA, nalidixic acid; OFX, ofloxacin; C, chloramphenicol; SD, sulfadiazine.

presence of IncF and IncI plasmids and was validated by PBRT. In addition, other plasmids were also identified by RT-PCR (*korB*, specific for IncP-1 plasmids) and Southern blot hybridization. Plasmids of known Inc groups were detected in all transconjugants from the mixed salad and cilantro but not in the transconjugants from arugula. Representative transconjugants from cilantro and mixed salad carried either one ( $n = 11$ ) or two ( $n = 3$ ) replicons. In 12 transconjugants from cilantro samples, four different plasmid replicon types were detected (Table 2), IncFII ( $n = 3$ ), IncFIB ( $n = 6$ ), IncI1 ( $n = 2$ ), and IncP-1β ( $n = 4$ ). In contrast, the transconjugants isolated from mixed salad showed only one replicon type, IncFII ( $n = 2$ ). One plasmid that could not be assigned by PBRT or RT-PCR was isolated from arugula leaves. The combination of replicon types IncFII and IncP-1β was detected in two transconjugants (pBC1.9 and pBC1.12), while the combination of replicon types of plasmids IncFIB and IncI1 was found in one transconjugant (pBC2.3) captured from cilantro leaves. Southern blot hybridization for sequences specific for IncP-1 plasmids revealed that four plasmids belonged to the IncP-1β subgroup. IncI2 plasmids were not detected in any TET-resistant transconjugants (Table 2). In contrast to IncFIB/FII and IncI1 plasmids, the IncP-1β plasmids captured exogenously were not detected in the 63 TET-resistant *E. coli* isolates.

#### Detection of IncF and IncI plasmids, *tet(A)*, and *int11* in total community DNA.

We also screened for plasmids (IncF, IncI1, and IncI2), tetracycline resistance gene *tet(A)*, and integrase gene *int11* in TC-DNA extracted from bacterial communities either directly after their detachment from fresh leaves or after an enrichment step, using PCR-Southern blot hybridization and RT-PCR (Table 3). Using the RT-PCR method, IncF and IncI plasmids as well as the *tet(A)* gene were detected in TC-DNA extracted from enrichment cultures of leaves, but not in TC-DNA from the detached bacteria. In contrast, the *int11* gene was detected in both kinds of TC-DNA. Consistent with these results, PCR-Southern blot hybridization targeting the IncF and IncI plasmids and *tet(A)* revealed strong hybridization signals in TC-DNA extracted from the enrichment cultures but very weak or no signals from direct extractions.

## DISCUSSION

The present study showed that bacteria associated with produce can carry various plasmids that might represent an important link between the environmental and the human gut microbiomes. Although initially low in abundance, TET-resistant *E. coli* were isolated from all purchased produce samples after nonselective enrichment. Contam-

**TABLE 3** PCR hybridization and real-time PCR of IncF, Inc11, and Inc12 plasmids and *int11* and *tet(A)* from TC-DNA extracted from produce before and after enrichment<sup>a</sup>

Produce	DNA isolation	Time point (day)	IncF		Inc11		Inc12		<i>int11</i>		<i>tet(A)</i>	
			RT-PCR	Blot	RT-PCR	Blot	RT-PCR	Blot	RT-PCR	Blot	RT-PCR	Blot
Mixed salad	Direct extraction	0	–	–	–	–	–	+++	+	(+++)	–	(++) <sup>1</sup>
		7	–	–	–	–	–	+++	+	(+++)	–	(+++) <sup>2</sup>
	Enrichment	0	+	(+++)	+	(++)	+	+++	+	(+++)	+	(++) <sup>2</sup>
		7	+	(+++)	+	(++)	+	+++	+	(+++)	+	+++
Arugula	Direct extraction	0	–	–	–	–	–	(++) <sup>3</sup>	+	(++)	–	(+++) <sup>2</sup>
		7	–	–	–	–	–	(++) <sup>1</sup>	+	(++)	–	(+++) <sup>1</sup>
	Enrichment	0	+	(+++)	–	–	+	+++	+	(++)	+	+++
		7	+	(+++)	–	–	+	+++	+	(++)	+	+++
Cilantro	Direct extraction	0	–	–	–	–	–	–	(+) <sup>1</sup>	(++)	(+) <sup>1</sup>	(+++) <sup>3</sup>
		7	(+) <sup>1</sup>	(++) <sup>2</sup>	–	–	–	–	(+) <sup>3</sup>	(+++)	(+) <sup>2</sup>	(+++) <sup>3</sup>
	Enrichment	0	(+) <sup>2</sup>	(+++) <sup>2</sup>	(+) <sup>1</sup>	(++) <sup>2</sup>	–	–	(+) <sup>3</sup>	(+++)	(+) <sup>2</sup>	(+++) <sup>3</sup>
		7	(+) <sup>2</sup>	(+++) <sup>2</sup>	(+) <sup>2</sup>	(++) <sup>2</sup>	–	–	(+) <sup>3</sup>	(+++) <sup>3</sup>	(+) <sup>3</sup>	(+++) <sup>3</sup>

<sup>a</sup>Superscript numbers indicate number of positive replicates; –, not detected or no signal; (+), positive (RT-PCR); (++) , medium signal; (+++), strong signal.

ination of produce with *E. coli* strains can occur in the field through contaminated soil (organic fertilizers), exposure to contaminated irrigation water, or during postharvest (12, 13). In this study, TET-resistant *E. coli* isolates were mostly isolated from cilantro that was purchased from supermarkets in Braunschweig and Magdeburg, Germany, followed by mixed ready-to-eat salad and arugula purchased from supermarkets in Braunschweig. This suggests that produce might be a hot spot for contamination with *E. coli* carrying multidrug resistance plasmids that occur at low abundance. A high proportion of the TET-resistant *E. coli* isolates was also resistant to penicillins (AM and AMX) and trimethoprim. Although it is difficult to compare among studies because of different methodologies used for isolation and resistance testing, our results are in line with high resistance levels to penicillins and trimethoprim previously reported for *E. coli* from irrigation water and vegetables (18), ready-to-eat salads (1), and lettuce (2). In the present study, TET resistance was commonly conferred by *tet(A)*, partly confirming previous studies reporting *tet(A)* and *tet(B)* genes as the most common TET resistance genes in *E. coli* and *Salmonella* spp. isolated from ready-to-eat vegetables (1, 33). The rapid dissemination of tetracycline resistance among bacteria has been related not only to the occurrence of TET resistance genes on transposons and conjugative plasmids (22, 23, 34), but also to selective pressure, e.g., the use of antibiotics in animal husbandry and the spread of TET resistance genes via organic fertilizers (31).

Plasmid-mediated multidrug resistance plays an important role in the transfer of ARGs around the world (35). Our study showed that *E. coli* isolates from produce harbored various plasmids belonging to replicon types IncF, Inc11, IncX1, IncU, IncN, and IncHI1, with IncF plasmids being the most frequently detected. These plasmids might play an important role in the dissemination of antibiotic resistances. IncF plasmids were found predominantly in *E. coli* isolated from drinking water (36) and poultry farms (37). In our study, IncFII was the most frequently detected replicon type (36.5%), followed by IncFIB (15.9%), which is in line with studies on *E. coli* recovered from pigs and humans (38), wastewater (39), and animals (40). The combination of replicon types IncFII and IncFIB in two isolates is consistent with a report on *Enterobacter cloacae* from lettuce (3). However, we cannot exclude that these replicons are located on the same plasmid, as several studies have reported the combination of replicon types as a multireplicon on a single plasmid (30, 41–43), likely due to cointegration (28). In this study, TET-resistant *E. coli* isolates which carried IncF plasmids were also positive for *tet(A)*, *aadA*, *sul1*, *sul2*, *sul3*, *qacE* and/or *qacEΔ1*, *qnrB*, *qnrS*, or *bla*<sub>TEM</sub> genes. Previous reports found that IncF plasmids can carry genes conferring resistance to all major antibiotic classes, including aminoglycosides, β-lactams, phenicols, tetracyclines, sulfonamides, and fluoroquinolones (38, 40, 44).

The NHR Inc11 plasmid types were the second most dominant replicon type (34.9%)

and IncI1-positive isolates also carried multiple ARGs. In this study, strains carrying IncI1 plasmids were also positive for class 1 integron integrase gene *intI1* and a diverse set of resistance genes, namely *tet(A)*, *sul2*, *strA*, *bla*<sub>TEM</sub>, *qacE* and/or *qacEΔ1*, *aadA*, *sul3*, *qnrS*, and/or *merRTΔP*. In a recent study, IncI1 plasmids from irrigation water and lettuce carried genes *sul1*, *tet(A)*, *aadA*, *strA*, and *bla*<sub>TEM</sub> as well as *intI1* (18). Similar phenotype and genotype profiles among *E. coli* strains from the current study and those recovered in previous studies from clinical samples, the environment, or other foods indicate that produce may play a potential role in the dispersal of *E. coli* carrying plasmid-localized ARGs. Thus, plasmids belonging to the IncF and IncI groups have the potential to be major contributors worldwide to the propagation of ARGs within enteric bacteria. One dissemination route of enteric bacteria carrying IncF and IncI plasmids might be the consumption of produce.

The newly developed TaqMan probe-based RT-PCR assays demonstrated high specificity in detecting these plasmids in *E. coli* isolates, and RT-PCR-positive isolates were also assigned by PBRT, which in addition enables subtyping.

This is the first study identifying NHR plasmids such as IncX1 and IncHI1 and broad-host-range (BHR) plasmid IncU in *E. coli* isolates recovered from cilantro leaves. Interestingly, IncX plasmids were detected in *E. cloacae* from lettuce (3). IncHI1 plasmids were previously reported in *E. coli* and *Citrobacter youngae* isolates from water and healthy calves, respectively (45), while the first IncU plasmids were isolated from *Aeromonas salmonicida* (46), and later from *Aeromonas caviae* from hospital effluent in the United Kingdom (47). In general, a low prevalence of ESBL-producing *E. coli* was found on produce, which is similar to previous studies (19, 48, 49). In the present study, ESBL-producing *E. coli* were isolated only from cilantro (2.8%).

To our knowledge, this is also the first report of *E. coli* isolates from cilantro that were positive for conjugative IncN plasmids, *bla*<sub>CTX-M-1</sub>, and resistance to third generation cephalosporins. The *bla*<sub>CTX-M-1</sub> gene was also reported on plasmids belonging to the IncN family in *E. coli* isolated from farm workers, animals, humans, and the environment (50–52). Although IncN plasmids are able to replicate in a variety of *Enterobacteriaceae*, they are most frequently found in *E. coli* and *Klebsiella pneumoniae*, where they contribute to the dissemination of cephalosporin and carbapenem resistance (53).

The results of the present study showed that *E. coli* isolates harboring the *bla*<sub>CTX-M-1</sub> gene also conferred resistance to at least seven classes of antibiotics tested. Moreover, *E. coli* harboring CTX-M genes were recently reported from lettuce and irrigation water (4, 54), raw vegetables (33, 54), and coastal waters (55, 56). Kim et al. (57) reported that ESBL-producing *E. coli* and *Klebsiella pneumoniae* carrying CTX-M were detected in ready-to-eat vegetables from a local retail market in Seoul, South Korea. A recent study has detected *bla*<sub>TEM</sub> genes in association with IncF and IncI1 plasmids from irrigation water and lettuce from 16 household farms in Estarreja, Portugal (18).

In previous reports, the occurrence of *sul1* and *qacEΔ1* was frequently associated with class 1 integrons (3, 58). Unexpectedly, only 27% and 36.5% of *sul1*- and *qacE/qacEΔ1*-positive isolates carried the *intI1* gene, respectively, indicating that atypical class 1 integrons were more prevalent among the isolates, as previously also reported by Amos et al. (59).

In the present study, transferable TET resistance plasmids were also directly captured from the produce microbiomes on day 0 but not on day 7 after purchase, and the highest transfer frequency was observed in mixed salad, followed by cilantro and arugula. Differences in observed frequencies of transconjugants could be due to different abundances of bacteria with conjugative plasmids in the various sample types, or due to real differences in the frequencies of plasmid transfer. The latter might be affected by the metabolic activity of the produce microbiome, as plasmid transfer frequency is known to depend not only on plasmid-specific characteristics, but also on ecological factors affecting the metabolic activity of bacteria (60). Replicon types IncFII, IncFIB, IncI1, and IncP-1β were captured from cilantro leaves, whereas only IncFII plasmids were captured from mixed salad. IncF (FII and FIB) plasmids were prevalent among TET-resistant transconjugants from both types of produce. Most of the IncF



plasmids exogenously captured harbored *bla*<sub>TEM</sub>, *tet(A)*, and *qnrS* genes. One IncI1 plasmid was captured from cilantro, and another one was captured in combination with replicon type IncFIB. The conjugative plasmids carried *tet(A)* and *bla*<sub>TEM</sub> genes. Finally, four IncP-1 $\beta$  plasmids were captured from cilantro leaves and two of them in combination with replicon type IncFII. IncP-1 plasmids have been frequently captured by exogenous plasmid isolation from various environments such as sewage sludge (61), manure (23), and water (62). However, the first isolations of IncP-1 plasmids were from clinical isolates (63, 64). The IncP-1 $\beta$  plasmids carried genes conferring resistances to antibiotics *tet(A)*, *strA*, and *bla*<sub>TEM</sub> and also mercury compounds (*merRT $\Delta$ P*) and disinfectants (*qacE/qacE $\Delta$ 1*).

In conclusion, this study showed that produce that we eat might contain bacteria such as *E. coli* carrying transferable multidrug resistance plasmids. Although *E. coli* numbers are typically low, our nonselective enrichments showed that proliferation can easily occur. Our study reports a specific TaqMan probe-based RT-PCR assay that can be used for rapid detection of IncF and IncI plasmids in *E. coli* isolates and exogenously captured plasmids as well as in TC-DNA extracted from enrichment cultures of leaves. However, quantifying these plasmids in TC-DNA directly extracted from the microbial fraction detached from leaves was impossible due to their low abundance in the microbiome, but IncF and IncI plasmids were detected in DNA extracted after previous enrichment. While these assays represent an important and useful tool to be implemented for monitoring the prevalence of IncF and IncI plasmids in isolates and the environment, negative results of these and other cultivation-independent methods can lead to an underestimation of the mobile resistome present in the rare microbiome of produce and other samples. This is the first study demonstrating that multidrug resistance plasmids present in produce-associated bacteria were transferable to sensitive *E. coli* recipients, a process that could occur in the human gut. The NHR plasmids IncF and IncI1 and also the BHR IncP-1 $\beta$  plasmids were captured from the produce. In particular, the captured IncF and IncI1 plasmids conferred resistance toward several classes of antibiotics. Thus, produce-associated bacteria should be considered an important route of disseminating transferable antibiotic resistances, which might be particularly relevant for patients under antibiotic treatment.

## MATERIALS AND METHODS

**Sample collection.** A total of 24 samples from different locally produced or imported produce (mixed salad, arugula, and cilantro) was analyzed. The mixed salad and arugula were purchased from local supermarkets in Braunschweig, Germany, in June and September 2016, and cilantro was obtained from supermarkets in Braunschweig and Magdeburg, Germany, in May 2017. The produce was stored at refrigerator temperature and sampled on days 0 and 7 (four replicates for each time point and produce type).

**Isolation and identification of TET-resistant *E. coli*.** For sampling, the produce was cut into pieces using a sterile scalpel and mixed. For each sample, 25 g each were filled in two stomacher bags (one for direct plating and the other for enrichment) and mixed three times with 75 ml buffered peptone water (BPW; Roth, Karlsruhe, Germany), with subsequent stomacher treatment performed with the Stomacher 400 (Seward, Worthing, United Kingdom) at high speed for 1 min. The enrichment cultures of fresh leaves in BPW were incubated at 37°C with shaking (150 rpm) for 18 to 24 h. In order to isolate TET-resistant *E. coli*, dilutions (10<sup>-1</sup> and 10<sup>-2</sup>) of the sample suspensions and 100  $\mu$ l of the enrichment cultures were plated on different culture media (eosin methylene blue [EMB]; Sifin, Berlin, Germany, and Chromocult coliform agar [CCA]; Merck, Darmstadt, Germany) supplemented with tetracycline (10 mg liter<sup>-1</sup>). All plates were incubated at 37°C for 18 to 24 h. The presumptive *E. coli* colonies were picked from each sample and streaked onto EMB, CCA, and TBX chromogenic agar (Roth, Karlsruhe, Germany) for confirmation by colony morphology and further characterization. *E. coli* isolates were then confirmed by biochemical tests for indole production, methyl red, and catalase activity (65). Furthermore, isolates were analyzed using PCR for the presence of the *gadA* gene encoding glutamate decarboxylase, specific for *E. coli* (66). *E. coli* isolates were stored in Luria broth (LB; Roth, Karlsruhe, Germany) containing 15% glycerol at -80°C.

**Exogenous plasmid isolation.** In order to capture tetracycline resistance plasmids, exogenous plasmid isolation via biparental mating was performed using *gfp*<sup>+</sup>, kanamycin (Km)<sup>-</sup>, and rifampin (Rif)<sup>-</sup> resistant *Escherichia coli* CV601 (67) as a recipient. The recipient strain was grown overnight in tryptic soy broth (TSB; Merck, Darmstadt, Germany) supplemented with rifampin (Rif) (50 mg liter<sup>-1</sup>) and kanamycin (Km) (50 mg liter<sup>-1</sup>). Two milliliters of the recipient strain culture was transferred into a sterile Eppendorf tube and centrifuged at 3,100  $\times$  *g* for 5 min and washed twice with 1:10 TSB. Then, the pellet was resuspended in 2 ml of 1:10 TSB. The bacterial suspensions (donor) of each sample on days 0 and

7 were prepared from enrichment cultures of fresh leaves as described above. Twenty milliliters of each enrichment culture (donor) and 0.5 ml of recipient strain were mixed in a 50-ml falcon tube. As a background control, 5 ml of the enrichment cultures and 200  $\mu$ l of the recipient were processed the same way as the samples. All mixtures were centrifuged at  $3,100 \times g$  for 10 min. The pellets were resuspended in 200  $\mu$ l of 1:10 TSB and then spotted onto a filter for mating (Millipore filters, 0.22  $\mu$ m). Filters were incubated overnight at 28°C on plate count agar plates (PCA; Merck, Darmstadt, Germany) supplemented with cycloheximide (Cyc) (100 mg liter<sup>-1</sup>). After incubation, the filters were placed in 2 ml of sterile 0.85% NaCl solution in a 50-ml falcon tube. Each filter was washed by vortexing for 1 min. Serial 10-fold dilutions were done and appropriate dilutions were plated on PCA agar supplemented with rifampin (Rif; 50 mg liter<sup>-1</sup>), kanamycin (Km; 50 mg liter<sup>-1</sup>), cycloheximide (Cyc; 100 mg liter<sup>-1</sup>), and tetracycline (TET; 15 mg liter<sup>-1</sup>) to select for tetracycline-resistant transconjugants. Background controls of bulk soil and the recipient controls were plated on the same selective media. Numbers of recipient cells were determined by applying three replicate 20- $\mu$ l drops per each serial dilution ( $10^{-5}$  to  $10^{-8}$ ) of all mating mixes on PCA with Km (50 mg liter<sup>-1</sup>), Rif (50 mg liter<sup>-1</sup>), and Cyc (100 mg liter<sup>-1</sup>). All plates were incubated at 28°C for up to 3 days. Transconjugants were determined by green fluorescence resulting from the green fluorescence protein (GFP). The identity of putative transconjugants was confirmed by BOX-PCR (68). Transfer frequencies were calculated as total number of transconjugants divided by the total number of recipients.

**Antibiotic susceptibility testing.** Antimicrobial susceptibility testing was performed by the disk diffusion method on Müller-Hinton agar (MH; Sigma-Aldrich, St. Louis, USA), according to the European Committee on Antimicrobial Susceptibility Testing (EUCAST). The antibiotics ( $\mu$ g) (Becton, Dickinson and Company, USA) used in this study were amoxicillin (25), ampicillin (10), cefotaxime (30), ceftazidime (30), ceftriaxone (30), chloramphenicol (30), ciprofloxacin (5), colistin (10), TET (30), doxycycline (30), streptomycin (10), gentamicin (10), ofloxacin (5), kanamycin (30), nalidixic acid (30), trimethoprim (5), and sulfadiazine (250). TET-resistant *E. coli* isolates were streaked onto LB agar supplemented with TET (10 mg liter<sup>-1</sup>), while TET-resistant *E. coli* CV601 transconjugants were streaked on plate count agar plates (PCA; Merck, Darmstadt, Germany) supplemented with TET (15 mg liter<sup>-1</sup>), Km (50 mg liter<sup>-1</sup>), and Rif (50 mg liter<sup>-1</sup>). *E. coli* strain CV601 was used as a negative control. The bacterial suspension was prepared from a single colony in normal saline (0.85% NaCl) to a density of 0.5 McFarland turbidity standard. Cotton swabs were used for streaking the suspension onto MH agar plates. After air drying, antibiotic discs were placed on the plates. Then all plates were incubated at 37°C for 18 to 24 h. The inhibition zone was measured. The results were interpreted according to the guidelines of EUCAST. Clinical and Laboratory Standards Institute (CLSI) recommendations were used when antibiotic breakpoints in EUCAST guidelines were absent (i.e., for doxycycline, streptomycin, tetracycline, and nalidixic acid). ESBL production was confirmed among TET-resistant *E. coli* isolates and transconjugants by double-disc diffusion test (DDT) (48). The ESBL producers were identified by phenotypic confirmatory test according to the CLSI.

**TC-DNA extraction.** The bacterial fraction detached from fresh leaves directly or after an enrichment culture of each sample as described above were pelleted by centrifugation at  $3,100 \times g$  for 15 min at 4°C. Total community DNA was extracted from the pellet using the FastDNA spin kit for soil (MP Biomedicals, Heidelberg, Germany), according to the manufacturer's instructions. The quality of extracted DNA was determined by agarose gel electrophoresis. The extracted DNA was stored at -20°C until further analysis.

**Genomic DNA extraction.** Genomic DNA was extracted from overnight cultures of TET-resistant *E. coli* isolates, transconjugants, and the recipient strain with a Qiagen genomic DNA extraction kit (Qiagen, Hilden, Germany) using a silica-based kit (silica bead DNA extraction kit; Thermo Scientific, St. Leon-Rot, Germany). The extracted genomic DNA was stored at -20°C until further analysis.

**Primer-probe design (IncF, Inc1, and Inc2 plasmids).** As it is known that relaxase genes can be used for classification of the mobilization systems of plasmids (69), the *tral* gene region was chosen as a target region to design primers detecting IncF, Inc1, and Inc2 plasmid sequences. A total of 4,530 plasmid DNA sequences were downloaded from NCBI (NCBI, Batch Entrez) using the 4,602 plasmid accession numbers found in GenBank by Shintani et al. (70), among which 298 plasmids were identified as belonging to the MOB<sub>F</sub> group. The coding sequences (CDS) of the MOB<sub>F</sub> plasmids were aligned using tBLASTn against the relaxase Tral of the F plasmid (GenBank accession number AP001918), resulting in 110 protein sequences sharing >50% identity and >70% coverage. The 110 protein sequences closely related to Tral were aligned using MAFT multiple sequence alignment software version 1.3.3. The alignment produced was back translated using the EMBOSS Backtranseq tool and used to generate a set of degenerated primers and probes using Primer3. All of those steps were carried out in Geneious 8.1.9. At best, 83 of the 110 *tral* nucleic acid sequences could be targeted by one set of designed primers and probe (Table 4). Those sequences belonged mostly to plasmids isolated from *Salmonella enterica* and *Escherichia coli* and a few from *Klebsiella pneumoniae* and *Shigella* spp. The plasmids corresponded to a part of the subclade MOB<sub>F12</sub> defined by Garcillán-Barcia et al. (71), which comprises the phylogenetically broad IncF complex. When tested against the 4,530 plasmids, the primer/probe targeted 92 plasmids in the database, and 89 of these plasmids belonged to the MOB<sub>F</sub> group (298 plasmids recovered belonged to this group) and three were annotated as "non-mob" and belonged to any MOB group. When looking at the "Inc" classification, 73 of the targeted plasmids belonged to the IncF (including FI and FII), 14 to the IncZ, and five to the pCD1 type. Among the 4,530 plasmid sequences, 243 carried a *rep* gene belonging to the IncF group, indicating that the primer/probe cannot detect all possible IncF plasmids.

**TABLE 4** PCR and RT-PCR primer systems used in this study

Gene target	Primer	Primer and probe (5'–3')	Size (bp)	Reference or source
<i>qacE</i> and/or <i>qacEΔ1</i>	qacEall-F	CGCATTATTTTCTTCTCTGGTT	69	76
	qacEall-R	CCCAGACCAGACTGCATAAGC		
	qacEall-P	TGAAATCCATCCCTGTCCGGTGT		
<i>tet(A)</i>	tetA-qfw	CCGCGCTTTGGGTCATT	504	77
	tetA_qrv	TGGTCGCGTCCCAGTGA		
	q-tetA-P	TCGGCGAGGATCG		
<i>sul1</i>	q-sul_1 653f	CCGTTGGCCTTCTGTAAAG	965	78
	q-sul_1 719r	TTGCCGATCGCGTGAAGT		
	tp_sul1	CAGCGAGCCTTGC GGCGG		
<i>sul2</i>	q_sul2 595f	CGGCTGCGCTTCGATT	865	78
	q_sul2 654f	CGCGCGCAGAAAGGATT		
	tp_sul2 614	CGGTGCTTCTGTCTGTTTCGGCG		
<i>sul3</i>	Sul3-F	CAGATAAGCAATTGAGCATGCTCTGC	569	38
	Sul3-R	AGAATGATTTCCGTGACACTGCAATCATT		
<i>intl1</i>	intl1-LC1	GCCTTGATGTTACCCGAGAG	196	79
	intl1-LC5	GATCGGTGGAATGCGTGT		
	intl1-P	ATTCCTGGCCGTGGTTCTGGGTTTT		
<i>aadA</i>	q-aadA-Fw	TTGATTTGCTGGTACTGTG	635	80
	q-aadA-Rv	CTTAGTGTGATCTCGCCTTT		
	q-aadA-P	TGGTAGGTCCAGCGCGGAG		
<i>korB</i>	korB-F	TCATCGACAACGACTACAACG	118	81
	korB-Fz	TCGTGGATAACGACTACAACG		
	korB-R	TTCTTCTGCCCTTCGCCAG		
	korB-Rge	TYTTCYTGCCCTTGCCAG		
	korB-Rd	TTCTTGACTCCCTTCGCCAG		
<i>strA</i>	q-strA-Fw	TCAATCCCGACTTCTTACCG	521	80
	q-strA-Rv	CACCATGGCAAACAACCATA		
	q-strA-P	TGCTCGACCAAGAGCGGC		
<i>intl2</i>	intl2-LC2	TGCTTTTCCCACCCTTACC	195	79
	intl2-LC3	GACGGCTACCCTCTGTTATCTC		
	intl2-P	TGGATACTCGCAACCAAGTTATTTTTACGCTG		
<i>tet(Q)</i>	q-tetQ-Fw	AGGTGCTGAACCTTGTTGATTC	69	82
	q-tetQ-Rv	GGCCGGACGGAGGATTT		
	q-tetQ-P	TCCGATCAGCATCCCCTC		
<i>IncF (tral)</i>	682_F	CACGGTATGTGGGARATGCC	391	This study
	1073_R	TCCGGCGGCAGYATVCCRAC		
	973_P	CAGCAGGGCGGTGRCRCAGGC		
<i>Inc11 (tral)</i>	Inc11_tral_Fwd	TTCTTCTCCCTACCATC	118	This study
	Inc11_tral_Rev	CATTTTCCAGCGTGTTTC		
	Inc11_tral_TP	CGGCTTTTCACTTCGTGGTT		
<i>Inc12 (tral)</i>	Inc12_tral_Fwd	CAAGAACAGAAACAGGCA	291	This study
	Inc12_tral_Rev	TCCCGCAGATAACAGATA		
	Inc12_tral_TP	CCAAACCAACCACAACCA		
<i>merRTΔP</i>	merRT-P	GGGAGATCTAAAGCACGCTAAGGCRTA	1000	83
	merRT-P	GGGGAATTCTTGACWGTGATCGGGCA		
<i>bla<sub>CTX-M-1</sub></i>	CTX-M-F	TCTTCCAGAATAAGGAATCCC	908	84
	CTX-M-R	CCGTTTCCGCTATTACAAC		
<i>bla<sub>TEM</sub></i>	TEM-F	TCCGCTCATGAGACAATAACC	930	84
	TEM-R	TTGGTCTGACAGTTACCAATGC		
<i>bla<sub>SHV</sub></i>	SHV-F	TTATCTCCCTGTTAGCCACC	796	84
	SHV-R	GATTTGCTGATTCGCTCGG		
<i>IS1071</i>	IS-F	GCTTGGTCACTTCTGGGTCTTC	180	85
	IS-R	CTATGCCCGTCTATCGTTACCC		
<i>IncN</i>	rep-1	AGTTCACCACCTACTCGCTCCG	165	86
	rep-2	CAAGTTCTTCTGTTGGGATTCGG		
<i>qnrA</i>	qnrAf-RT	ATTTCTCACGCCAGGATTTG	529	87
<i>qnrB</i>	qnrAr-RT	GCAGATCGGCATAGCTGAAG	429	87
	qnrBmF	GGMATHGAAATTCGCCACTG		
<i>qnrS</i>	qnrBmR	TTYGCBYYCGCCAGTCGAA	393	87
	qnrSrtF11	GACGTGCTAACTTGCGTGAT		
	qnrSrtR11	TGGCATTGTTGGAAACTTG		

Available IncI plasmid sequences were downloaded from NCBI. The *tral* genes were aligned using the software CLC Main Workbench version 8 (CLC bio, Qiagen) with standard settings for alignments, and primers were designed to match conserved regions of the *tral* gene (Table 4). The specificity was confirmed *in silico* with NCBI primer BLAST and with a set of plasmids from other incompatibility groups. Plasmids used for this test were R388, pB10, pHHV216, RSF1010, pSM1890, RP4, pHH3-414, pHH2-227,

pRA3, RN3, RSF1010, pTH10, pTP6, R751, pQKH54, pKS208, pEST4002, pJKJ5, pMCBF1, pRMS149, pCAR1, pD2RT, pD67, pWW0, and pST527, from which none was amplified.

**Detection of IncF, IncI plasmids by real-time PCR.** The RT-PCR assay was performed under standard conditions. All real-time PCR (RT-PCR) reactions were set up in a 25  $\mu$ l reaction volume using a Hot Start *Taq* DNA polymerase (M0495L, New England BioLabs, Ipswich, Massachusetts, USA) containing 5  $\mu$ l of template DNA, 300 nM primer (reverse), 50 nM primer (forward), and 50 nM probe for IncF, and 300 nM primer each (forward and reverse) and TaqMan probe for IncI. All primers and probes are described in Table 4. The following PCR program was used for amplification: 10 min at 95°C, followed by 40 cycles of 95°C for 30 sec and 60°C for 1 min. The assays were carried out in triplicate with real-time PCR 5'-nuclease assays (TaqMan RT-PCR) in a CFX96 real-time PCR detection system (Bio-Rad, Hercules, CA, USA). Negative controls were included in all tests, and they consisted of all the elements of the reaction except for the template DNA.

Standard plasmids were used to construct a full standard curve in duplicate in each RT-PCR run. Standard plasmids were constructed by cloning the purified PCR products amplified from the plasmids R64 for IncI1, pHNSHP45 for IncI2, and F plasmid for IncF using the corresponding primer pairs used for the RT-PCR, into TransforMax EC100 electrocompetent *E. coli* (Epicentre), in pJET1.2 using the Thermo Scientific CloneJET PCR cloning kit (Thermo Scientific).

**Detection of target genes by real-time PCR and PCR.** The target genes in genomic DNA extracted from TET-resistant *E. coli* isolates and transconjugants were detected by RT-PCR 5'-nuclease assays (TaqMan or EvaGreen RT-PCR) in a CFX96 RT-PCR detection system (Bio-Rad, Hercules, CA, USA) or by PCR for class 1 and 2 integrons; integrase genes *int1* and *int2*; *korB* (IncP-1 plasmids); *qacE* and/or *qacE $\Delta$ 1* (*qacE/qacE $\Delta$ 1*) encoding quaternary ammonium compound resistance; *aadA* and *strA* encoding streptomycin and spectinomycin resistance; tetracycline resistance genes [*tet(Q)* and *tet(A)*]; the *merRT $\Delta$ P* gene part of the mercury resistance operon; *sul1*, *sul2*, and *sul3* encoding sulfonamide resistance; *qnrA*, *qnrB*, and *qnrS* encoding fluoroginolone resistance;  $\beta$ -lactam resistance genes (*bla<sub>TEM</sub>*, *bla<sub>SHV</sub>* and *bla<sub>CTX-M-1</sub>*); IncN (*rep*); and insertion sequence (IS) *IS1071* (represented by the *tnpA* gene). The DNA of the recipient strain *E. coli* CV601 was included as the negative control. The primers and probes targeting these genes and PCR and RT-PCR conditions are listed in Table 4.

**Conjugation assay.** TET-resistant *E. coli* isolates positive for ESBL (EK2.29, EK3.43, and EK3.44) were examined for their ability to transfer resistance. Briefly, the donors and the rifampin- and kanamycin-resistant *E. coli* CV601 recipient strain were grown in LB broth overnight at 37°C. 500  $\mu$ l of overnight cultures of each donor and recipient strain were mixed in 1 ml LB broth and incubated at 37°C for 24 h without shaking. 100  $\mu$ l of the conjugal mixture was spread on LB agar (Roth; Karlsruhe, Germany) containing Rif (50 mg liter<sup>-1</sup>), Kan (50 mg liter<sup>-1</sup>), CTX (2 mg liter<sup>-1</sup>), and TET (15 mg liter<sup>-1</sup>), and incubated at 37°C for 24 to 48 h. The transconjugants were verified by BOX-PCR and further tested for the antibiotic resistance phenotypes and genotypes as described above.

**Plasmid DNA extraction and detection by Southern blot hybridization.** Plasmid DNA extraction from TET-resistant transconjugants (pBC1.1, pBC1.3, pBC1.9, and pBC1.12) captured exogenously from the enrichment of cilantro leaves was performed using Qiagen Plasmid Mini Kit (Qiagen, Inc., Hilden, Germany) according to the manufacturer's instructions. In order to detect the IncP-1 plasmids with Southern blot hybridization in these transconjugants, plasmid DNA was digested with the restriction enzyme NotI (Thermo Fisher Scientific, Waltham, MA, USA), and fragments were separated by electrophoresis on a 1% agarose-TBE gel as described previously (72). Southern blot hybridization was performed with digoxigenin (DIG)-labeled probes generated from PCR amplicons which were obtained from reference plasmid R751 for IncP1- $\beta$  as previously described by Binh et al. (73).

**Plasmid replicon typing.** PBRT was used to identify the incompatibility group of plasmids in TET-resistant *E. coli* and transconjugants, and to confirm the presence of IncF and IncI plasmids as determined via the newly developed RT-PCR method as described above. This was done by PCR amplification on genomic DNA of the strains using primer sets for 30 replicons, HI1, HI2, I1, I2, X1, X2, X3, X4, L, M, N, FIA, FIB, FIC, FII, FIIS, FIIK, FIB KN, FIB KQ, W, Y, P1, A/C, T, K, U, R, B/O, HIB-M, and FIB-M, representative of major plasmid incompatibility groups among *Enterobacteriaceae* (74, 28). PCR products were separated by electrophoresis on a 2.5% agarose-TBE gel and stained with ethidium bromide.

**Detection of IncF and IncI plasmids, *tet(A)*, and the class 1 integrase gene *int1* via PCR-Southern blot hybridization and RT-PCR in TC-DNA.** PCR-Southern blot hybridization or RT-PCR was used to detect *tet(A)*, *int1*, IncF, and IncI plasmids in TC-DNA extracted from the microbial fraction detached from leaves directly or after an enrichment step on days 0 and 7. The PCR products were separated on a 1% agarose-TBE gel electrophoresis and then transferred to a positively charged nylon membrane (GE Healthcare, UK). Southern blot hybridization was carried out with digoxigenin (DIG)-labeled probes generated from PCR amplicons, which were obtained from reference plasmids pJKJ5 for *int1* and *tet(A)*, as described by Dealtry et al. (75), R64 for IncI1, and pHNSHP45 for IncI2 and IncF. The primers and PCR conditions are listed in Table 4.

## ACKNOWLEDGMENTS

We thank Ute Zimmerling and Lena Rauch for their technical assistance.

K.B. had a scholarship from the Libyan government, S.J. was funded by the German Environment Agency (Umweltbundesamt) (FKZ 3713 63402), and T.S. was supported by the Agriculture and Food Research Initiative competitive grant 2013-67019-21375 from the USDA National Institute of Food and Agriculture.

## REFERENCES

- Campos J, Mourão J, Pestana N, Peixe L, Novais C, Antunes P. 2013. Microbiological quality of ready-to-eat salads: an underestimated vehicle of bacteria and clinically relevant antibiotic resistance genes. *Int J Food Microbiol* 166:464–470. <https://doi.org/10.1016/j.ijfoodmicro.2013.08.005>.
- Holvoet K, Sampers I, Callens B, Dewulf J, Uyttendaele M. 2013. Moderate prevalence of antimicrobial resistance in *Escherichia coli* isolates from lettuce, irrigation water, and soil. *Appl Environ Microbiol* 79:6677–6683. <https://doi.org/10.1128/AEM.01995-13>.
- Jones-Dias D, Manageiro V, Ferreira E, Barreiro P, Vieira L, Moura IB, Caniça M. 2016. Architecture of Class 1, 2, and 3 integrons from Gram negative bacteria recovered among fruits and vegetables. *Front Microbiol* 7:1400. <https://doi.org/10.3389/fmicb.2016.01400>.
- Njage PMK, Buys EM. 2015. Pathogenic and commensal *Escherichia coli* from irrigation water show potential in transmission of extended spectrum and AmpC  $\beta$ -lactamases determinants to isolates from lettuce. *Microb Biotechnol* 8:462–473. <https://doi.org/10.1111/1751-7915.12234>.
- Buchholz U, Bernard H, Werber D, Böhmer MM, Remschmidt C, Wilking H, Deleré Y, An der Heiden M, Adlhoeh C, Dreesman J, Ehlers J, Ethelberg S, Faber M, Frank C, Fricke G, Greiner M, Höhle M, Ivarsson S, Jark U, Kirchner M, Koch J, Krause G, Lubber P, Rosner B, Stark K, Kühne M. 2011. German outbreak of *Escherichia coli* O104:H4 associated with sprouts. *N Engl J Med* 365:1763–1770. <https://doi.org/10.1056/NEJMoa1106482>.
- Callejón RM, Rodríguez-Naranjo MI, Ubeda C, Hornedo-Ortega R, Garcia-Parrilla MC, Troncoso AM. 2015. Reported foodborne outbreaks due to fresh produce in the United States and European Union: trends and causes. *Foodborne Pathog Dis* 12:32–38. <https://doi.org/10.1089/fpd.2014.1821>.
- Herman KM, Hall AJ, Gould LH. 2015. Outbreaks attributed to fresh leafy vegetables, United States, 1973–2012. *Epidemiol Infect* 143:3011–3021. <https://doi.org/10.1017/S0950268815000047>.
- Wendel AM, Johnson DH, Sharapov U, Grant J, Archer JR, Monson T, Koschmann C, Davis JP. 2009. Multistate outbreak of *Escherichia coli* O157:H7 infection associated with consumption of packaged spinach, August–September 2006: the Wisconsin investigation. *Clin Infect Dis* 48:1079–1086. <https://doi.org/10.1086/597399>.
- Friesema I, Sigmundsdottir G, van der Zwaluw K, Heuvelink A, Schimmer B, de Jager C, Rupp B, Briem H, Hardardottir H, Atladottir A, Gudmundsdottir E, van Pelt W. 2008. An international outbreak of Shiga toxin-producing *Escherichia coli* O157 infection due to lettuce, September–October 2007. *Eurosurveillance* 13:19065. <https://doi.org/10.2807/ese.13.50.19065-en>.
- Berg G, Grube M, Schlöter M, Smalla K. 2014. The plant microbiome and its importance for plant and human health. *Front Microbiol* 5:491. <https://doi.org/10.3389/fmicb.2014.00491>.
- Sáenz Y, Zarazaga M, Briñas L, Lantero M, Ruiz-Larrea F, Torres C. 2001. Antibiotic resistance in *Escherichia coli* isolates obtained from animals, foods and humans in Spain. *Int J Antimicrob Agents* 18:353–358. [https://doi.org/10.1016/S0924-8579\(01\)00422-8](https://doi.org/10.1016/S0924-8579(01)00422-8).
- Berger CN, Sodha SV, Shaw RK, Griffin PM, Pink D, Hand P, Frankel G. 2010. Fresh fruit and vegetables as vehicles for the transmission of human pathogens. *Environ Microbiol* 12:2385–2397. <https://doi.org/10.1111/j.1462-2920.2010.02297.x>.
- Alegbeleye OO, Singleton I, Sant'Ana AS. 2018. Sources and contamination routes of microbial pathogens to fresh produce during field cultivation. A review. *Food Microbiol* 73:177–208. <https://doi.org/10.1016/j.fm.2018.01.003>.
- Gustafson RH, Bowen RE. 1997. Antibiotic use in animal agriculture. *J Appl Microbiol* 83:531–541. <https://doi.org/10.1046/j.1365-2672.1997.00280.x>.
- McManus PS, Stockwell VO, Sundin GW, Jones AL. 2002. Antibiotic use in plant agriculture. *Annu Rev Phytopathol* 40:443–465. <https://doi.org/10.1146/annurev.phyto.40.120301.093927>.
- Chang Q, Wang W, Regev-Yochay G, Lipsitch M, Hanage WP. 2015. Antibiotics in agriculture and the risk to human health. How worried should we be? *Evol Appl* 8:240–247. <https://doi.org/10.1111/eva.12185>.
- Grenni P, Ancona V, Barra Caracciolo A. 2018. Ecological effects of antibiotics on natural ecosystems: a review. *Microchemical J* 136:25–39. <https://doi.org/10.1016/j.microc.2017.02.006>.
- Araújo S, Silva IAT, Tação M, Patinha C, Alves A, Henriques I. 2017. Characterization of antibiotic resistant and pathogenic *Escherichia coli* in irrigation water and vegetables in household farms. *Int J Food Microbiol* 257:192–200. <https://doi.org/10.1016/j.ijfoodmicro.2017.06.020>.
- Blaak H, van Hoek AHAM, Veenman C, Docters van Leeuwen AE, Lynch G, van Overbeek WM, de Roda Husman AM. 2014. Extended spectrum  $\beta$ -lactamase- and constitutively AmpC-producing Enterobacteriaceae on fresh produce and in the agricultural environment. *Int J Food Microbiol* 168:169:8–16. <https://doi.org/10.1016/j.ijfoodmicro.2013.10.006>.
- Hudson JA, Frewer LJ, Jones G, Brereton PA, Whittingham MJ, Stewart G. 2017. The agri-food chain and antimicrobial resistance. A review. *Trends Food Sci Technol* 69:131–147. <https://doi.org/10.1016/j.tifs.2017.09.007>.
- van Elsas JD, Turner S, Bailey MJ. 2003. Horizontal gene transfer in the phytosphere. *New Phytol* 157:525–537. <https://doi.org/10.1046/j.1469-8137.2003.00697.x>.
- Bennett PM. 2009. Plasmid encoded antibiotic resistance: acquisition and transfer of antibiotic resistance genes in bacteria. *Br J Pharmacol* 153:5347–5357. <https://doi.org/10.1038/sj.bjpp.0707607>.
- Heuer H, Binh CTT, Jechalke S, Kopmann C, Zimmerling U, Krögerrecklenfort E, Ledger T, González B, Top E, Smalla K. 2012. IncP-1 $\epsilon$  plasmids are important vectors of antibiotic resistance genes in agricultural systems: diversification driven by class 1 Integron gene cassettes. *Front Microbiol* 3:2. <https://doi.org/10.3389/fmicb.2012.00002>.
- Verraes C, van Boxtael S, van Meerveen E, van Coillie E, Butaye P, Catry B, Schaezen M-A, de van Huffel X, Imberechts H, Dierick K, Daube G, Saegerman C, Block J, de Dewulf J, Herman L. 2013. Antimicrobial resistance in the food chain: a review. *Int J Environ Res Public Health* 10:2643–2669. <https://doi.org/10.3390/ijerph10072643>.
- Song J, Rensing C, Holm PE, Virta M, Brandt KK. 2017. Comparison of metals and tetracycline as selective agents for development of tetracycline resistant bacterial communities in agricultural soil. *Environ Sci Technol* 51:3040–3047. <https://doi.org/10.1021/acs.est.6b05342>.
- Suzuki H, Yano H, Brown CJ, Top EM. 2010. Predicting plasmid promiscuity based on genomic signature. *J Bacteriol* 192:6045–6055. <https://doi.org/10.1128/JB.00277-10>.
- Rozwandowicz M, Brouwer MSM, Fischer J, Wagenaar JA, Gonzalez-Zorn B, Guerra B, Mevius DJ, Hordijk J. 2018. Plasmids carrying antimicrobial resistance genes in *Enterobacteriaceae*. *J Antimicrob Chemother* 73:1121–1137. <https://doi.org/10.1093/jac/dkx488>.
- Villa L, García-Fernández A, Fortini D, Carattoli A. 2010. Replicon sequence typing of IncF plasmids carrying virulence and resistance determinants. *J Antimicrob Chemother* 65:2518–2529. <https://doi.org/10.1093/jac/dkq347>.
- Johnson TJ, Nolan LK. 2009. Pathogenomics of the virulence plasmids of *Escherichia coli*. *Microbiol Mol Biol Rev* 73:750–774. <https://doi.org/10.1128/MMBR.00015-09>.
- Koraimann G. 2018. Spread and persistence of virulence and antibiotic resistance genes: a ride on the F plasmid conjugation module. *EcoSal Plus* 8. <https://doi.org/10.1128/ecosalplus.ESP-0003-2018>.
- Wolters B, Widyasari-Mehta A, Kreuzig R, Smalla K. 2016. Contaminations of organic fertilizers with antibiotic residues, resistance genes, and mobile genetic elements mirroring antibiotic use in livestock? *Appl Microbiol Biotechnol* 100:9343–9353. <https://doi.org/10.1007/s00253-016-7742-y>.
- Smalla K, Jechalke S, Top EM. 2015. Plasmid detection, characterization and ecology. *Microbiol Spectr* 3. <https://doi.org/10.1128/microbiolspec.PLAS-0038-2014>.
- Skočková A, Karpíšková R, Koláčková I, Cupáková Š. 2013. Characteristics of *Escherichia coli* from raw vegetables at a retail market in the Czech Republic. *Int J Food Microbiol* 167:196–201. <https://doi.org/10.1016/j.ijfoodmicro.2013.09.011>.
- Jones-Dias D, Manageiro V, Caniça M. 2016. Influence of agricultural practice on mobile *bla* genes: Inc11-bearing CTX-M, SHV, CMY and TEM in *Escherichia coli* from intensive farming soils. *Environ Microbiol* 18:260–272. <https://doi.org/10.1111/1462-2920.13021>.
- Wang J, Stephan R, Karczmarczyk M, Yan Q, Hächler H, Fanning S. 2013. Molecular characterization of *bla* ESBL-harboring conjugative plasmids identified in multi-drug resistant *Escherichia coli* isolated from food-producing animals and healthy humans. *Front Microbiol* 4:188. <https://doi.org/10.3389/fmicb.2013.00188>.
- Lyimo B, Buza J, Subbiah M, Temba S, Kipasika H, Smith W, Call DR. 2016. IncF plasmids are commonly carried by antibiotic resistant *Escherichia*

- coli* isolated from drinking water sources in Northern Tanzania. Int J Microbiol 2016:1. <https://doi.org/10.1155/2016/3103672>.
37. Johnson TJ, Wannemuehler YM, Johnson SJ, Logue CM, White DG, Doetkott C, Nolan LK. 2007. Plasmid replicon typing of commensal and pathogenic *Escherichia coli* isolates. Appl Environ Microbiol 73: 1976–1983. <https://doi.org/10.1128/AEM.02171-06>.
  38. Wu S, Dalsgaard A, Hammerum AM, Porsbo LJ, Jensen LB. 2010. Prevalence and characterization of plasmids carrying sulfonamide resistance genes among *Escherichia coli* from pigs, pig carcasses and human. Acta Vet Scand 52:47. <https://doi.org/10.1186/1751-0147-52-47>.
  39. Wibberg D, Szczepanowski R, Eikmeyer F, Pühler A, Schlüter A. 2013. The IncF plasmid pRSB225 isolated from a municipal wastewater treatment plant's on-site preflooder combining antibiotic resistance and putative virulence functions is highly related to virulence plasmids identified in pathogenic *E. coli* isolates. Plasmid 69:127–137. <https://doi.org/10.1016/j.plasmid.2012.11.001>.
  40. Yang Q-E, Sun J, Li L, Deng H, Liu B-T, Fang L-X, Liao X-P, Liu Y-H. 2015. IncF plasmid diversity in multi-drug resistant *Escherichia coli* strains from animals in China. Front Microbiol 6:964. <https://doi.org/10.3389/fmicb.2015.00964>.
  41. Carattoli A. 2009. Resistance plasmid families in *Enterobacteriaceae*. Antimicrob Agents Chemother 53:2227–2238. <https://doi.org/10.1128/AAC.01707-08>.
  42. Bengtsson S, Naseer U, Sundsfjord A, Kahlmeter G, Sundqvist M. 2012. Sequence types and plasmid carriage of uropathogenic *Escherichia coli* devoid of phenotypically detectable resistance. J Antimicrob Chemother 67:69–73. <https://doi.org/10.1093/jac/dkr421>.
  43. Osborn AM, da Silva Tatley FM, Steyn LM, Pickup RW, Saunders JR. 2000. Mosaic plasmids and mosaic replicons: evolutionary lessons from the analysis of genetic diversity in IncFII-related replicons. Microbiol 146: 2267–2275. <https://doi.org/10.1099/00221287-146-9-2267>.
  44. Liu B-T, Yang Q-E, Li L, Sun J, Liao X-P, Fang L-X, Yang S-S, Deng H, Liu Y-H. 2013. Dissemination and characterization of plasmids carrying *oqxAB-bla<sub>CTX-M</sub>* genes in *Escherichia coli* isolates from food-producing animals. PLoS One 8:e73947. <https://doi.org/10.1371/journal.pone.0073947>.
  45. Zurfluh K, Jakobi G, Stephan R, Hächler H, Nüesch-Inderbinen M. 2014. Replicon typing of plasmids carrying *bla<sub>CTX-M-1</sub>* in *Enterobacteriaceae* of animal, environmental and human origin. Front Microbiol 5:555. <https://doi.org/10.3389/fmicb.2014.00555>.
  46. Aoki T, Mitoma Y, Crosa JH. 1986. The characterization of a conjugative R-plasmid isolated from *Aeromonas salmonicida*. Plasmid 16:213–218. [https://doi.org/10.1016/0147-619X\(86\)90059-4](https://doi.org/10.1016/0147-619X(86)90059-4).
  47. Rhodes G, Parkhill J, Bird C, Ambrose K, Jones MC, Huys G, Swings J, Pickup RW. 2004. Complete nucleotide sequence of the conjugative tetracycline resistance plasmid pFBAOT6, a member of a group of IncU plasmids with global ubiquity. Appl Environ Microbiol 70:7497–7510. <https://doi.org/10.1128/AEM.70.12.7497-7510.2004>.
  48. Bhutani N, Muraleedharan C, Talreja D, Rana SW, Walia S, Kumar A, Walia SK. 2015. Occurrence of multidrug resistant extended spectrum beta-lactamase-producing bacteria on iceberg lettuce retailed for human consumption. Biomed Res Int 2015:1. <https://doi.org/10.1155/2015/547547>.
  49. Raphael E, Wong LK, Riley LW. 2011. Extended-spectrum beta-lactamase gene sequences in Gram-negative saprophytes on retail organic and nonorganic spinach. Appl Environ Microbiol 77:1601–1607. <https://doi.org/10.1128/AEM.02506-10>.
  50. Dolejska M, Villa L, Hasman H, Hansen L, Carattoli A. 2013. Characterization of IncN plasmids carrying *bla<sub>CTX-M-1</sub>* and *qnr* genes in *Escherichia coli* and *Salmonella* from animals, the environment and humans. J Antimicrob Chemother 68:333–339. <https://doi.org/10.1093/jac/dks387>.
  51. Moodley A, Guardabassi L. 2009. Transmission of IncN plasmids carrying *bla<sub>CTX-M-1</sub>* between commensal *Escherichia coli* in pigs and farm workers. Antimicrob Agents Chemother 53:1709–1711. <https://doi.org/10.1128/AAC.01014-08>.
  52. Novais A, Cantón R, Moreira R, Peixe L, Baquero F, Coque TM. 2007. Emergence and dissemination of *Enterobacteriaceae* isolates producing CTX-M-1-like enzymes in Spain are associated with IncFII (CTX-M-15) and broad-host-range (CTX-M-1, -3, and -32) plasmids. Antimicrob Agents Chemother 51:796–799. <https://doi.org/10.1128/AAC.01070-06>.
  53. Carattoli A. 2013. Plasmids and the spread of resistance. Int J Med Microbiol 303:298–304. <https://doi.org/10.1016/j.ijmm.2013.02.001>.
  54. Ben Said L, Jouini A, Klibi N, Dziri R, Alonso CA, Boudabous A, Ben Slama K, Torres C. 2015. Detection of extended-spectrum beta-lactamase (ESBL)-producing *Enterobacteriaceae* in vegetables, soil and water of the farm environment in Tunisia. Int J Food Microbiol 203:86–92. <https://doi.org/10.1016/j.ijfoodmicro.2015.02.023>.
  55. Leonard AFC, Zhang L, Balfour AJ, Garside R, Hawkey PM, Murray AK, Ukoumunne OC, Gaze WH. 2018. Exposure to and colonisation by antibiotic-resistant *E. coli* in UK coastal water users: environmental surveillance, exposure assessment, and epidemiological study (Beach Bum Survey). Environ Int 114:326–333. <https://doi.org/10.1016/j.envint.2017.11.003>.
  56. Alves MS, Pereira A, Araújo SM, Castro BB, Correia ACM, Henriques I. 2014. Seawater is a reservoir of multi-resistant *Escherichia coli*, including strains hosting plasmid-mediated quinolones resistance and extended-spectrum beta-lactamases genes. Front Microbiol 5:426. <https://doi.org/10.3389/fmicb.2014.00426>.
  57. Kim H-S, Chon J-W, Kim Y-J, Kim M-s, Seo K-H. 2015. Prevalence and characterization of extended-spectrum-β-lactamase-producing *Escherichia coli* and *Klebsiella pneumoniae* in ready-to-eat vegetables. Int J Food Microbiol 207:83–86. <https://doi.org/10.1016/j.ijfoodmicro.2015.04.049>.
  58. Gillings MR, Gaze WH, Pruden A, Smalla K, Tiedje JM, Zhu Y-G. 2015. Using the class 1 integron-integrase gene as a proxy for anthropogenic pollution. ISME J 9:1269–1279. <https://doi.org/10.1038/ismej.2014.226>.
  59. Amos GCA, Ploumaki S, Zhang L, Hawkey PM, Gaze WH, Wellington EMH. 2018. The widespread dissemination of integrons throughout bacterial communities in a riverine system. ISME J 12:681–691. <https://doi.org/10.1038/s41396-017-0030-8>.
  60. van Elsas JD, Bailey MJ. 2002. The ecology of transfer of mobile genetic elements. FEMS Microbiol Ecol 42:187–197. <https://doi.org/10.1111/j.1574-6941.2002.tb01008.x>.
  61. Schlüter A, Szczepanowski R, Pühler A, Top EM. 2007. Genomics of IncP-1 antibiotic resistance plasmids isolated from wastewater treatment plants provides evidence for a widely accessible drug resistance gene pool. FEMS Microbiol Rev 31:449–477. <https://doi.org/10.1111/j.1574-6976.2007.00074.x>.
  62. Brown CJ, Sen D, Yano H, Bauer ML, Rogers LM, van der Auwera GA, Top EM. 2013. Diverse broad-host-range plasmids from freshwater carry few accessory genes. Appl Environ Microbiol 79:7684–7695. <https://doi.org/10.1128/AEM.02252-13>.
  63. Thorsted PB, Macartney DP, Akhtar P, Haines AS, Ali N, Davidson P, Stafford T, Pocklington MJ, Pansegrau W, Wilkins BM, Lanka E, Thomas CM. 1998. Complete sequence of the IncP(beta) plasmid R751: implications for evolution and organisation of the IncP backbone. J Mol Biol 282:969–990. <https://doi.org/10.1006/jmbi.1998.2060>.
  64. Thomas CM, Smith CA. 1987. Incompatibility group P plasmids: Genetics, evolution, and use in genetic manipulation. Annu Rev Microbiol 41: 77–101. <https://doi.org/10.1146/annurev.mi.41.100187.000453>.
  65. Harley JP, Prescott LM. 2002. Laboratory exercises in microbiology, 5th ed. McGraw-Hill, Boston, MA.
  66. McDaniels AE, Rice EW, Reyes AL, Johnson CH, Haugland RA, Stelma GN. 1996. Confirmational identification of *Escherichia coli*, a comparison of genotypic and phenotypic assays for glutamate decarboxylase and β-D-glucuronidase. Appl Environ Microbiol 62:3350–3354.
  67. Heuer H, Krögerrecklenfort E, Wellington EMH, Egan S, Elsas JD, Overbeek L, Collard J-M, Guillaume G, Karagouni AD, Nikolakopoulou TL, Smalla K. 2002. Gentamicin resistance genes in environmental bacteria: prevalence and transfer. FEMS Microbiol Ecol 42:289–302. <https://doi.org/10.1111/j.1574-6941.2002.tb01019.x>.
  68. Rademaker LW, Frank J, Frans JB. 1997. Characterization of the diversity of ecologically important microbes by rep-PCR genomic fingerprinting, p.1–26. In Akkermans ADL, van Elsas JD, de Bruin FJ (ed), Molecular microbial ecology manual II, 2nd ed. Kluwer Academic Publishers, Dordrecht, The Netherlands.
  69. Alvarado A, Garcillán-Barcia MP, de La Cruz F. 2012. A degenerate primer MOB typing (DPMT) method to classify gamma-proteobacterial plasmids in clinical and environmental settings. PLoS One 7:e40438. <https://doi.org/10.1371/journal.pone.0040438>.
  70. Shintani M, Sanchez ZK, Kimbara K. 2015. Genomics of microbial plasmids: classification and identification based on replication and transfer systems and host taxonomy. Front Microbiol 6:242. <https://doi.org/10.3389/fmicb.2015.00242>.
  71. Garcillán-Barcia MP, Francia MV, de La Cruz F. 2009. The diversity of conjugative relaxases and its application in plasmid classification. FEMS Microbiol Rev 33:657–687. <https://doi.org/10.1111/j.1574-6976.2009.00168.x>.

72. Sambrook J, Fritsch EF, Maniatis T (ed). 1989. Molecular cloning: a laboratory manual, 2nd ed. Cold Spring Harbor Laboratory Press, New York, NY.
73. Binh CTT, Heuer H, Kaupenjohann M, Smalla K. 2008. Piggery manure used for soil fertilization is a reservoir for transferable antibiotic resistance plasmids. *FEMS Microbiol Ecol* 66:25–37. <https://doi.org/10.1111/j.1574-6941.2008.00526.x>.
74. Carattoli A, Bertini A, Villa L, Falbo V, Hopkins KL, Threlfall EJ. 2005. Identification of plasmids by PCR-based replicon typing. *J Microbiol Methods* 63:219–228. <https://doi.org/10.1016/j.mimet.2005.03.018>.
75. Dealtry S, Ding G-C, Weichelt V, Dunon V, Schlüter A, Martini MC, Del Papa MF, Lagares A, Amos GCA, Wellington EMH, Gaze WH, Sipkema D, Sjöling S, Springael D, Heuer H, van Elsas JD, Thomas C, Smalla K. 2014. Cultivation-independent screening revealed hot spots of IncP-1, IncP-7 and IncP-9 plasmid occurrence in different environmental habitats. *PLoS One* 9:e89922. <https://doi.org/10.1371/journal.pone.0089922>.
76. Jechalke S, Schreiter S, Wolters B, Dealtry S, Heuer H, Smalla K. 2013. Widespread dissemination of class 1 integron components in soils and related ecosystems as revealed by cultivation-independent analysis. *Front Microbiol* 4:420. <https://doi.org/10.3389/fmicb.2013.00420>.
77. Guarddon M, Miranda JM, Rodríguez JA, Vázquez BI, Cepeda A, Franco CM. 2011. Real-time polymerase chain reaction for the quantitative detection of *tetA* and *tetB* bacterial tetracycline resistance genes in food. *Int J Food Microbiol* 146:284–289. <https://doi.org/10.1016/j.ijfoodmicro.2011.02.026>.
78. Heuer H, Smalla K. 2007. Manure and sulfadiazine synergistically increased bacterial antibiotic resistance in soil over at least two months. *Environ Microbiol* 9:657–666. <https://doi.org/10.1111/j.1462-2920.2006.01185.x>.
79. Barraud O, Baclet MC, Denis F, Ploy MC. 2010. Quantitative multiplex real-time PCR for detecting class 1, 2 and 3 integrons. *J Antimicrob Chemother* 65:1642–1645. <https://doi.org/10.1093/jac/dkq167>.
80. Walsh F, Ingenfeld A, Zampiccoli M, Hilber-Bodmer M, Frey JE, Duffy B. 2011. Real-time PCR methods for quantitative monitoring of streptomycin and tetracycline resistance genes in agricultural ecosystems. *J Microbiol Methods* 86:150–155. <https://doi.org/10.1016/j.mimet.2011.04.011>.
81. Jechalke S, Dealtry S, Smalla K, Heuer H. 2013. Quantification of IncP-1 plasmid prevalence in environmental samples. *Appl Environ Microbiol* 79:1410–1413. <https://doi.org/10.1128/AEM.03728-12>.
82. Smith MS, Yang RK, Knapp CW, Niu Y, Peak N, Hanfelt MM, Galland JC, Graham DW. 2004. Quantification of tetracycline resistance genes in feedlot lagoons by real-time PCR. *Appl Environ Microbiol* 70:7372–7377. <https://doi.org/10.1128/AEM.70.12.7372-7377.2004>.
83. Osborn AM, Bruce KD, Strike P, Ritchie DA. 1993. Polymerase chain reaction-restriction fragment length polymorphism analysis shows divergence among *mer* determinants from Gram-negative soil bacteria indistinguishable by DNA-DNA hybridization. *Appl Environ Microbiol* 59:4024–4030.
84. Stürenburg E, Kühn A, Mack D, Laufs R. 2004. A novel extended-spectrum beta-lactamase CTX-M-23 with a P167T substitution in the active-site omega loop associated with ceftazidime resistance. *J Antimicrob Chemother* 54:406–409. <https://doi.org/10.1093/jac/dkh334>.
85. Providenti MA, O'Brien JM, Ewing RJ, Paterson ES, Smith ML. 2006. The copy-number of plasmids and other genetic elements can be determined by SYBR-Green-based quantitative real-time PCR. *J Microbiol Methods* 65:476–487. <https://doi.org/10.1016/j.mimet.2005.09.007>.
86. Götz A, Pukall R, Smit E, Tietze E, Prager R, Tschäpe H, van Elsas JD, Smalla K. 1996. Detection and characterization of broad-host-range plasmids in environmental bacteria by PCR. *Appl Environ Microbiol* 62:2621–2628.
87. Marti E, Balcázar JL. 2013. Real-time PCR assays for quantification of *qnr* genes in environmental water samples and chicken feces. *Appl Environ Microbiol* 79:1743–1745. <https://doi.org/10.1128/AEM.03409-12>.